

## Vector Valued multiple of $\chi^2$ over $p$ -metric sequence spaces defined by Musielak

Vandana<sup>1</sup>, Deepmala<sup>2</sup>, N. Subramanian<sup>3</sup> and Lakshmi Narayan Mishra<sup>4</sup>

<sup>1</sup> Department of Management Studies, Indian Institute of Technology, Madras, Chennai 600 036, Tamil Nadu, India. email: [vdrai1988@gmail.com](mailto:vdrai1988@gmail.com)

<sup>2</sup> Mathematics Discipline, PDPM Indian Institute of Information Technology, Design and Manufacturing, Jabalpur, P.O.: Khamaria, Jabalpur 482 005, Madhya Pradesh, India. , email: [dmrai23@gmail.com](mailto:dmrai23@gmail.com)

<sup>3</sup> Department of Mathematics, SASTRA University, Thanjavur-613 401, India, email: [nsmaths@yahoo.com](mailto:nsmaths@yahoo.com)

<sup>4</sup> Department of Mathematics, Lovely Professional University, Jalandhar-Delhi G.T. Road, Phagwara, Punjab 144 411, India, L. 1627 Awadh Puri Colony Beniganj, Phase -III, Opposite-Industrial Training Institute (I.T.I.), Ayodhya Main Road Faizabad 224 001, Uttar Pradesh, India email: [lakshminarayanmishra04@gmail.com](mailto:lakshminarayanmishra04@gmail.com), [l\\_n\\_mishra@yahoo.co.in](mailto:l_n_mishra@yahoo.co.in)

**ABSTRACT.** In this article, we define the vector valued multiple of  $\chi^2$  over  $p$ - metric sequence spaces defined by Musielak and study some of their topological properties and some inclusion results.

**Keywords:** Analytic sequence, double sequences,  $\chi^2$  space, Musielak - modulus function,  $p$ - metric space, multiplier space.

*2000 Mathematics subject classification:* 40A05; 40C05; Secondary 46A45; 03E72.

---

<sup>1</sup> Corresponding author: [lakshminarayanmishra04@gmail.com](mailto:lakshminarayanmishra04@gmail.com)  
Received: 6 February 2016  
Revised: 21 September 2016  
Accepted: 26 October 2016

## 1. INTRODUCTION

Consider  $w, \chi$  and  $\Lambda$  denote the classes of all, gai and analytic scalar valued single sequences, respectively. We write  $w^2$  for the set of all complex sequences  $(x_{mn})$ , where  $m, n \in \mathbb{N}$ , the set of positive integers. Then,  $w^2$  is a linear space under the coordinate wise addition and scalar multiplication. Throughout this article the space of regularly gai multiple sequence defined over a semi-normed space  $(X, q)$ , semi-normed by  $q$  will be denoted by  $\chi_{mn}^{2R}(q)$  and  $\Lambda_{mn}^{2R}(q)$  For  $X = \mathbb{C}$ , the field of complex numbers, these spaces represent the corresponding scalar sequence spaces. Some initial works on double sequence spaces is found in Bromwich [1]. Later on, they were investigated by Hardy [2], Moricz [3], Moricz and Rhoades [4], Basarir and Solankan [5], Tripathy [6], Turkmenoglu [7], and many others. We procure the following sets of double sequences:

$$\begin{aligned} \mathcal{M}_u(t) &:= \{(x_{mn}) \in w^2 : \sup_{m,n \in \mathbb{N}} |x_{mn}|^{t_{mn}} < \infty\}, \\ \mathcal{C}_p(t) &:= \\ \{(x_{mn}) \in w^2 : p\text{-}\lim_{m,n \rightarrow \infty} |x_{mn} - l|^{t_{mn}} = 1 \text{ for some } l \in \mathbb{C}\}, \\ \mathcal{C}_{0p}(t) &:= \{(x_{mn}) \in w^2 : p\text{-}\lim_{m,n \rightarrow \infty} |x_{mn}|^{t_{mn}} = 1\}, \\ \mathcal{L}_u(t) &:= \{(x_{mn}) \in w^2 : \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} |x_{mn}|^{t_{mn}} < \infty\}, \\ \mathcal{C}_{bp}(t) &:= \mathcal{C}_p(t) \cap \mathcal{M}_u(t) \text{ and } \mathcal{C}_{0bp}(t) = \mathcal{C}_{0p}(t) \cap \mathcal{M}_u(t); \end{aligned}$$

where  $t = (t_{mn})$  is the sequence of strictly positive reals  $t_{mn}$  for all  $m, n \in \mathbb{N}$  and  $p\text{-}\lim_{m,n \rightarrow \infty}$  denotes the limit in the Pringsheim's sense. In the case  $t_{mn} = 1$  for all  $m, n \in \mathbb{N}$ ;  $\mathcal{M}_u(t), \mathcal{C}_p(t), \mathcal{C}_{0p}(t), \mathcal{L}_u(t), \mathcal{C}_{bp}(t)$  and  $\mathcal{C}_{0bp}(t)$  reduce to the sets  $\mathcal{M}_u, \mathcal{C}_p, \mathcal{C}_{0p}, \mathcal{L}_u, \mathcal{C}_{bp}$  and  $\mathcal{C}_{0bp}$ , respectively. Now, we may summarize the knowledge given in some document related to the double sequence spaces. Gökhan and Colak [8,9] have proved that  $\mathcal{M}_u(t)$  and  $\mathcal{C}_p(t), \mathcal{C}_{bp}(t)$  are complete paranormed spaces of double sequences and gave the  $\alpha-, \beta-, \gamma-$  duals of the spaces  $\mathcal{M}_u(t)$  and  $\mathcal{C}_{bp}(t)$ . Quite recently, in her PhD thesis, Zelter [10] has essentially studied both the theory of topological double sequence spaces and the theory of summability of double sequences. Mursaleen and Edely [11] and Tripathy have independently introduced the statistical convergence and Cauchy for double sequences and given the relation between statistical convergent and strongly Cesàro summable double sequences. Altay and Basar [12] have defined the spaces  $\mathcal{BS}, \mathcal{BS}(t), \mathcal{CS}_p, \mathcal{CS}_{bp}, \mathcal{CS}_r$  and  $\mathcal{BV}$  of double sequences consisting of all double series whose sequence of partial sums are in the spaces  $\mathcal{M}_u, \mathcal{M}_u(t), \mathcal{C}_p, \mathcal{C}_{bp}, \mathcal{C}_r$  and  $\mathcal{L}_u$ , respectively, and also examined some properties of those sequence spaces and determined the  $\alpha-$  duals of the spaces  $\mathcal{BS}, \mathcal{BV}, \mathcal{CS}_{bp}$  and the  $\beta(\vartheta)-$  duals of the spaces  $\mathcal{CS}_{bp}$  and  $\mathcal{CS}_r$  of double series. Basar and Sever [13]

have introduced the Banach space  $\mathcal{L}_q$  of double sequences corresponding to the well-known space  $\ell_q$  of single sequences and examined some properties of the space  $\mathcal{L}_q$ . Quite recently Subramanian and Misra [14] have studied the space  $\chi_M^2(p, q, u)$  of double sequences and gave some inclusion relations.

The class of sequences which are strongly Cesàro summable with respect to a modulus was introduced by Maddox [15] as an extension of the definition of strongly Cesàro summable sequences. Connor [16] further extended this definition to a definition of strong  $A$ -summability with respect to a modulus where  $A = (a_{n,k})$  is a nonnegative regular matrix and established some connections between strong  $A$ -summability, strong  $A$ -summability with respect to a modulus, and  $A$ -statistical convergence. In [17] the notion of convergence of double sequences was presented by A. Pringsheim. Also, in [18]-[19], and [20] the four dimensional matrix transformation  $(Ax)_{k,\ell} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{k\ell}^{mn} x_{mn}$  was studied extensively by Robison and Hamilton.

We need the following inequality in the sequel of the paper. For  $a, b, \geq 0$  and  $0 < p < 1$ , we have

$$(a + b)^p \leq a^p + b^p \quad (1.1)$$

The double series  $\sum_{m,n=1}^{\infty} x_{mn}$  is called convergent if and only if the double sequence  $(s_{mn})$  is convergent, where  $s_{mn} = \sum_{i,j=1}^{m,n} x_{ij}$  ( $m, n \in \mathbb{N}$ ).

A sequence  $x = (x_{mn})$  is said to be double analytic if,  $\sup_{m,n} |x_{mn}|^{1/m+n} < \infty$ . The vector space of all double analytic sequences will be denoted by  $\Lambda^2$ . A sequence  $x = (x_{mn})$  is called double gai sequence if  $((m+n)! |x_{mn}|)^{1/m+n} \rightarrow 0$  as  $m, n \rightarrow \infty$ . The double gai sequences will be denoted by  $\chi^2$ . Let  $\phi = \{finite\ sequences\}$ . Consider a double sequence  $x = (x_{ij})$ . The  $(m, n)^{th}$  section  $x^{[m,n]}$  of the sequence is defined by  $x^{[m,n]} = \sum_{i,j=0}^{m,n} x_{ij} \mathfrak{S}_{ij}$  for all  $m, n \in \mathbb{N}$ ; where  $\mathfrak{S}_{ij}$  denotes the double sequence whose only non zero term is a  $\frac{1}{(i+j)!}$  in the  $(i, j)^{th}$  place for each  $i, j \in \mathbb{N}$ .

An FK-space (or a metric space)  $X$  is said to have AK property if  $(\mathfrak{S}_{mn})$  is a Schauder basis for  $X$ . Or equivalently  $x^{[m,n]} \rightarrow x$ .

t An FDK-space is a double sequence space endowed with a complete metrizable; locally convex topology under which the coordinate mappings  $x = (x_k) \rightarrow (x_{mn})$  ( $m, n \in \mathbb{N}$ ) are also continuous.

Let  $M$  and  $\Phi$  are mutually complementary modulus functions. Then, we have:

(i) For all  $u, y \geq 0$ ,

$$uy \leq M(u) + \Phi(y), (Young's\ inequality)[See[21]] \quad (1.2)$$

(ii) For all  $u \geq 0$ ,

$$u\eta(u) = M(u) + \Phi(\eta(u)). \quad (1.3)$$

(iii) For all  $u \geq 0$ , and  $0 < \lambda < 1$ ,

$$M(\lambda u) \leq \lambda M(u) \quad (1.4)$$

Lindenstrauss and Tzafriri [22] used the idea of Orlicz function to construct Orlicz sequence space

$$\ell_M = \left\{ x \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty, \text{ for some } \rho > 0 \right\},$$

The space  $\ell_M$  with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\},$$

becomes a Banach space which is called an Orlicz sequence space. For  $M(t) = t^p$  ( $1 \leq p < \infty$ ), the spaces  $\ell_M$  coincide with the classical sequence space  $\ell_p$ .

A sequence  $f = (f_{mn})$  of modulus function is called a Musielak-modulus function. A sequence  $g = (g_{mn})$  defined by

$$g_{mn}(v) = \sup \{ |v|u - (f_{mn})(u) : u \geq 0 \}, m, n = 1, 2, \dots$$

is called the complementary function of a Musielak-modulus function  $f$ . For a given Musielak modulus function  $f$ , the Musielak-modulus sequence space  $t_f$  is defined as follows

$$t_f = \left\{ x \in w^2 : M_f(|x_{mn}|)^{1/m+n} \rightarrow 0 \text{ as } m, n \rightarrow \infty \right\},$$

where  $M_f$  is a convex modular defined by

$$M_f(x) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} f_{mn}(|x_{mn}|)^{1/m+n}, x = (x_{mn}) \in t_f.$$

We consider  $t_f$  equipped with the Luxemburg metric

$$d(x, y) = \sup_{mn} \left\{ \inf \left( \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} f_{mn} \left( \frac{|x_{mn} - y_{mn}|}{mn} \right)^{1/m+n} \right) \leq 1 \right\}$$

If  $X$  is a sequence space, we give the following definitions:

(i)  $X'$  = the continuous dual of  $X$ ;

(ii)  $X^\alpha = \{ a = (a_{mn}) : \sum_{m,n=1}^{\infty} |a_{mn}x_{mn}| < \infty, \text{ for each } x \in X \}$ ;

(iii)  $X^\beta = \{ a = (a_{mn}) : \sum_{m,n=1}^{\infty} a_{mn}x_{mn} \text{ is convergent, for each } x \in X \}$ ;

(iv)  $X^\gamma = \left\{ a = (a_{mn}) : \sup_{mn \geq 1} \left| \sum_{m,n=1}^{M,N} a_{mn}x_{mn} \right| < \infty, \text{ for each } x \in X \right\}$ ;

(v) let  $X$  be an  $FK$ -space  $\supset \phi$ ; then  $X^f = \{ f(\mathfrak{S}_{mn}) : f \in X' \}$ ;

$$(vi) X^\delta = \left\{ a = (a_{mn}) : \sup_{mn} |a_{mn} x_{mn}|^{1/m+n} < \infty, \text{ for each } x \in X \right\};$$

$X^\alpha, X^\beta, X^\gamma$  are called  $\alpha$ - (or Köthe-Toeplitz) dual of  $X, \beta$ - (or generalized-Köthe-Toeplitz) dual of  $X, \gamma$ - dual of  $X, \delta$ - dual of  $X$  respectively.  $X^\alpha$  is defined by Gupta and Kamphan . It is clear that  $X^\alpha \subset X^\beta$  and  $X^\alpha \subset X^\gamma$ , but  $X^\beta \subset X^\gamma$  does not hold, since the sequence of partial sums of a double convergent series need not to be bounded.

The notion of difference sequence spaces (for single sequences) was introduced by Kizmaz as follows

$$Z(\Delta) = \{x = (x_k) \in w : (\Delta x_k) \in Z\}$$

for  $Z = c, c_0$  and  $\ell_\infty$ , where  $\Delta x_k = x_k - x_{k+1}$  for all  $k \in \mathbb{N}$ .

Here  $c, c_0$  and  $\ell_\infty$  denote the classes of convergent, null and bounded scalar valued single sequences respectively. The difference sequence space  $bv_p$  of the classical space  $\ell_p$  is introduced and studied in the case  $1 \leq p \leq \infty$  by Başar and Altay and in the case  $0 < p < 1$  by Altay and Başar. The spaces  $c(\Delta), c_0(\Delta), \ell_\infty(\Delta)$  and  $bv_p$  are Banach spaces normed by

$$\|x\| = |x_1| + \sup_{k \geq 1} |\Delta x_k| \text{ and } \|x\|_{bv_p} = (\sum_{k=1}^{\infty} |x_k|^p)^{1/p}, (1 \leq p < \infty).$$

Later on the notion was further investigated by many others. We now introduce the following difference double sequence spaces defined by

$$Z(\Delta) = \{x = (x_{mn}) \in w^2 : (\Delta x_{mn}) \in Z\}$$

where  $Z = \Lambda^2, \chi^2$  and  $\Delta x_{mn} = (x_{mn} - x_{mn+1}) - (x_{m+1n} - x_{m+1n+1}) = x_{mn} - x_{mn+1} - x_{m+1n} + x_{m+1n+1}$  for all  $m, n \in \mathbb{N}$ .

## 2. DEFINITION AND PRELIMINARIES

Throughout this article a multiple sequence is denoted by;

$A = \langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle$ , a multiple infinite array of elements

$a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \in X$  for all  $m_1 m_2 \dots m_r n_1 n_2 \dots n_s \in \mathbb{N}$ .

Let  $n \in \mathbb{N}$  and  $X$  be a real vector space of dimension  $m$ , where  $n \leq m$ .

A real valued function  $d_p(x_1, \dots, x_n) = \|(d_1(x_1), \dots, d_n(x_n))\|_p$  on  $X$  satisfying the following four conditions:

(i)  $\|(d_1(x_1), \dots, d_n(x_n))\|_p = 0$  if and only if  $d_1(x_1), \dots, d_n(x_n)$  are linearly dependent,

(ii)  $\|(d_1(x_1), \dots, d_n(x_n))\|_p$  is invariant under permutation,

(iii)  $\|(\alpha_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} d_1(x_1), \dots, \alpha_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} d_n(x_n))\|_p = |\alpha_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s}| \|(d_1(x_1), \dots, d_n(x_n))\|_p,$

$\alpha_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \in \mathbb{R}$

(iv)  $d_p((x_1, y_1), (x_2, y_2) \dots (x_n, y_n)) = (d_X(x_1, x_2, \dots, x_n))^p + d_Y(y_1, y_2, \dots, y_n)^p)^{1/p}$  for  $1 \leq p < \infty$ ; (or)

(v)  $d((x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)) := \sup \{d_X(x_1, x_2, \dots, x_n), d_Y(y_1, y_2, \dots, y_n)\}$ , for  $x_1, x_2, \dots, x_n \in X, y_1, y_2, \dots, y_n \in Y$  is called the  $p$  product metric of

the Cartesian product of  $n$  metric spaces is the  $p$  norm of the  $n$ -vector of the norms of the  $n$  subspaces.

A trivial example of  $p$  product metric of  $n$  metric space is the  $p$  norm space is  $X = \mathbb{R}$  equipped with the following Euclidean metric in the product space is the  $p$  norm:

$$\|(d_1(x_1), \dots, d_n(x_n))\|_E = \sup (|\det(d_{mn}(x_{mn}))|) = \sup \left( \begin{array}{cccc} d_{11}(x_{11}) & d_{12}(x_{12}) & \dots & d_{1n}(x_{1n}) \\ d_{21}(x_{21}) & d_{22}(x_{22}) & \dots & d_{2n}(x_{2n}) \\ \vdots & \vdots & \ddots & \vdots \\ d_{n1}(x_{n1}) & d_{n2}(x_{n2}) & \dots & d_{nn}(x_{nn}) \end{array} \right)$$

where  $x_i = (x_{i1}, \dots, x_{in}) \in \mathbb{R}^n$  for each  $i = 1, 2, \dots, n$ .

If every Cauchy sequence in  $X$  converges to some  $L \in X$ , then  $X$  is said to be complete with respect to the  $p$ - metric. Any complete  $p$ - metric space is said to be  $p$ - Banach metric space.

**2.1. Definition.** Let  $X$  be a linear metric space. A function  $\rho : X \rightarrow \mathbb{R}$  is called paranorm, if

- (1)  $\rho(x) \geq 0$ , for all  $x \in X$ ;
- (2)  $\rho(-x) = \rho(x)$ , for all  $x \in X$ ;
- (3)  $\rho(x + y) \leq \rho(x) + \rho(y)$ , for all  $x, y \in X$ ;
- (4) If  $(\sigma_{mn})$  is a sequence of scalars with  $\sigma_{mn} \rightarrow \sigma$  as  $m, n \rightarrow \infty$  and  $(x_{mn})$  is a sequence of vectors with  $\rho(x_{mn} - x) \rightarrow 0$  as  $m, n \rightarrow \infty$ , then  $\rho(\sigma_{mn}x_{mn} - \sigma x) \rightarrow 0$  as  $m, n \rightarrow \infty$ .

A paranorm  $w$  for which  $\rho(x) = 0$  implies  $x = 0$  is called total paranorm and the pair  $(X, w)$  is called a total paranormed space. It is well known that the metric of any linear metric space is given by some total paranorm (see [23], Theorem 10.4.2, p.183).

**2.2. Definition.** A multiple sequence space  $E$  is said to be solid if

$\langle \alpha_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in E$  whenever  $\langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in E$  for all multiple sequences  $\langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle$  of scalars with  $|\alpha_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s}| \leq 1$  for all  $m_1 m_2 \dots m_r n_1 n_2 \dots n_s \in \mathbb{N}$ .

**2.3. Definition.** A multiple sequence space  $E$  is said to be symmetric if

$\langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in E$ , implies  $\langle a_{\pi(m_1 m_2 \dots m_r n_1 n_2 \dots n_s)} \rangle \in E$ ,

where  $\pi(m_1 m_2 \dots m_r n_1 n_2 \dots n_s) \in E$  are permutations of  $(M \times N) \dots (M \times N)$ .

**2.4. Definition.** A multiple sequence space  $E$  is said to be monotone if it contains the canonical pre-images of all its step spaces.

**2.5. Defintion.** A multiple sequence space  $E$  is said to be convergence free if  $\langle b_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in E$ , whenever  $\langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in E$  and  $b_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} = 0$  whenever  $a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} = 0$ .

**2.6. Remark.** A sequence space  $E$  is solid implies  $E$  is monotone.

Let  $f$  be an Musielak modulus function. Now we introduce the following multiple sequence spaces:

$$\left[ \Lambda_{mn}^{2fq}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] = \langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in w_{mn}^{2q} :$$

$$\left[ f \left( q \left( \eta_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] < \infty,$$

$$\text{where } \eta_{uv}(x) = |a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s}|^{(1/m_1 m_2 \dots m_r) + n_1 n_2 \dots n_s}$$

$$\left[ \chi_{mn}^{2fq}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] = \langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in w_{mn}^{2q} :$$

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \rightarrow 0;$$

$$as m_1 m_2 \dots m_r n_1 n_2 \dots n_s \rightarrow \infty.$$

where  $\mu_{uv}(x) = ((m_1 m_2 \dots m_r + n_1 n_2 \dots n_s)! |a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s}|)^{(1/m_1 m_2 \dots m_r) + n_1 n_2 \dots n_s}$   
 $A = \langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in \chi_{mn}^{2Rf}(q)$ , (i.e.), regularly gai if  $\langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in \chi_{mn}^{2f}(q)$  and the following limit hold:

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \rightarrow 0; as m_1 n_1 \rightarrow \infty$$

and  $m_2 \dots m_r, n_2 \dots n_s \in \mathbb{N}$

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \rightarrow 0;$$

$as m_2 n_2 \rightarrow \infty$  and  $m_3 \dots m_r, n_3 \dots n_s \in \mathbb{N}$

$\vdots$   
 $\vdots$

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \rightarrow 0 as m_r n_s \rightarrow \infty$$

and  $m_1 \dots m_{r-1}, n_1 \dots n_{s-1} \in \mathbb{N}$

**2.7. Remark.** The space  $\chi_{mn}^{2Rf}(q)$  has the following definition too

$$\left[ \chi_{mn}^{2fq}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] = \langle a_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} \rangle \in w_{mn}^{2q} :$$

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \rightarrow 0;$$

$$as max \{m_1 m_2 \dots m_r n_1 n_2 \dots n_s\} \rightarrow \infty. \chi_{mn}^{2Bf}(q) = \chi_{mn}^{2f}(q) \cap \Lambda_{mn}^{2f}(q).$$

### 3. MAIN RESULT

**Theorem 3.1.**  $\chi_{mn}^{2Rf}(q)$ ,  $\chi_{mn}^{2Bf}(q)$  and  $\Lambda_{mn}^{2f}(q)$  of multiple sequences are linear spaces.

**Proof:** We have to use linearity condition and then prove to the statement. Hence it is trivial. Therefore omit the proof.

**Theorem 3.2.** Let  $f = (f_{mn})$  be a multiple sequence of Musielak-modulus functions. Then the space

$\left[ \chi_{mn}^{2Rf}(q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  is a paranormed space with respect to the paranorm defined by

$$g(x) = \inf \left\{ \left( \left[ f \left( q \left( \|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \right\} \leq 1.$$

**Proof:** Clearly  $g(x) \geq 0$  for  $x = (a_{m_1 m_2 \dots m_r, n_1 n_2 \dots n_s}) \in \left[ \chi_{mn}^{2Rf}(q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$ . Since  $f(0) = 0$ , we get  $g(0) = 0$ .

Conversely, suppose that  $g(x) = 0$ , then

$$\inf \left\{ \left( \left[ f \left( q \left( \|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \right\} \leq 1 = 0.$$

Suppose that  $\mu_{uv}(x) \neq 0$  for each  $u, v \in \mathbb{N}$ . Then

$\|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \rightarrow \infty$ . It follows that

$\left( \left[ f \left( q \left( \|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \rightarrow \infty$  which is a contradiction. Therefore  $\mu_{uv}(x) = 0$ . Let

$$\left( \left[ f \left( q \left( \|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1 \text{ and}$$

$$\left( \left[ f \left( q \left( \|\mu_{uv}(y), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1.$$

Then by using Minkowski's inequality, we have

$$\begin{aligned} & \left( \left[ f \left( q \left( \|\mu_{uv}(x+y), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq \\ & \left( \left[ f \left( q \left( \|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) + \\ & \left( \left[ f \left( q \left( \|\mu_{uv}(y), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right). \end{aligned}$$

So we have  $g(x+y) = \inf$

$$\begin{aligned} & \left\{ \left( \left[ f \left( q \left( \|\mu_{uv}(x+y), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1 \right\} \leq \\ & \inf \left\{ \left( \left[ f \left( q \left( \|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1 \right\} + \\ & \inf \left\{ \left( \left[ f \left( q \left( \|\mu_{uv}(y), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1 \right\} \end{aligned}$$

Therefore,

$$g(x+y) \leq g(x) + g(y).$$

Finally, to prove that the scalar multiplication is continuous. Let  $\lambda_{m_1 m_2 \dots m_r, n_1 n_2 \dots n_s}$  be any complex number. By definition,

$$g(\lambda_{m_1 m_2 \dots m_r, n_1 n_2 \dots n_s} x) = \inf$$



$$\left\{ \left( \left[ f \left( q \left( \|\mu_{uv}(\lambda_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1 \right\}$$

Then

$$g(\lambda_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} x) = \inf$$

$$\left\{ \left( (\lambda_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} | t) : \left( \left[ f \left( q \left( \|\mu_{uv}(x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1 \right) \right\}$$

where  $t = \frac{1}{|\lambda|}$ . Since  $|\lambda| \leq \max(1, |\lambda|^{supp_{uv}})$ , we have

$$g(\lambda_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} x) \leq \max(1, |\lambda|^{supp_{uv}})$$

$$\inf \left\{ t : \left( \left[ f \left( q \left( \|\mu_{uv}(\lambda_{m_1 m_2 \dots m_r n_1 n_2 \dots n_s} x), (d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \right) \leq 1 \right\}.$$

This completes the proof.

**Theorem 3.3.** The space  $\left[ \chi_{mn}^{2Rf}(q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  is not symmetric.

**Proof:** Let  $\langle a_{m_1 m_2 \dots m_r, n_1 n_2 \dots n_s} \rangle \in \left[ \chi_{mn}^{2Rf}(q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$ .

Then for a given  $\epsilon > 0$  there exists a positive integers  $g_1, g_2, \dots, g_{g+1} h_1, h_2 \dots h_{h+1}$  such that

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] < \epsilon \text{ for all } m_1 n_1 > g_1 h_1 \text{ for all } m_2 \dots m_r, n_2 \dots n_s \in \mathbb{N}$$

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] < \epsilon \text{ for all } m_2 n_2 > g_2 h_2 \text{ for all } m_3 \dots m_r, n_3 \dots n_s \in \mathbb{N}$$

$\vdots$   
 $\vdots$

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] < \epsilon \text{ for all } m_r n_s > g_r h_s \text{ for all } m_1 \dots m_{r-1}, n_1 \dots n_{s-1} \in \mathbb{N}$$

$$\left[ f \left( q \left( \mu_{uv}(x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] < \epsilon \text{ for all } m_1 > g_{r+1}, m_2 > g_{r+1}, \dots, m_r > g_{r+1}, n_1 > h_{s+1}, n_2 > h_{s+1}, \dots, n_s > h_{s+1} \in \mathbb{N}.$$

Let  $g_0 h_0 = \max \{g_1, g_2, \dots, g_r, g_{r+1}, h_1, h_2 \dots h_r, h_{r+1}\}$ .

Let  $\langle b_{m_1 m_2 \dots m_r, n_1 n_2 \dots n_s} \rangle$  be a rearrangement of  $\langle a_{m_1 m_2 \dots m_r, n_1 n_2 \dots n_s} \rangle$ . Then

we have  $a_{i_1 i_2 \dots i_r, j_1 j_2 \dots j_s} = b_{m_{i_1} m_{i_2} \dots m_{i_r}, n_{j_1} n_{j_2} \dots n_{j_s}}$  for all  $i_1 i_2 \dots i_r, j_1 j_2 \dots j_s \in \mathbb{N}$ . Let

$$g_{r+2} h_{r+2} = \max$$

$$m_1 m_2 \dots m_{r_1}, m_{r+1}, m_{(r_0)_1} n_1 n_2 \dots n_{s_1}, n_{s+1} \dots m_{1_r} m_{2_r} \dots m_{r_r}, m_{r+1_r}, m_{(r_0)_r} n_{1_s} n_{2_s} \dots n_{s_s}, n_{s+1_s}.$$

Then we have

$\left[ f \left( q \left( \mu_{uv} (x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] < \epsilon$  for all  $m_1 n_1 > g_{r+2} h_{s+2} \cdots m_r n_s > g_{r+2} h_{s+2}$ .  
 Thus  $\left\langle b_{m_{i_1} m_{i_2} \cdots m_{i_r}, n_{j_1} n_{j_2} \cdots n_{j_s}} \right\rangle \in \left[ \chi_{mn}^{2Rf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$ .  
 Hence  $\left[ \chi_{mn}^{2Rf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  is a symmetric space.

**Theorem 3.4.** *The spaces  $\left[ \chi_{mn}^{2Rf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  and  $\left[ \chi_{mn}^{2Bf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  are solid*

**Proof:** The spaces  $\left[ \chi_{mn}^{2Rf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  and  $\left[ \chi_{mn}^{2Bf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  are solid follows the following inequality.

$\left[ f \left( q \left( \alpha_{m_1 m_2 \cdots m_r, n_1 n_2 \cdots n_s} \mu_{uv} (x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right] \leq \left[ f \left( q \left( \mu_{uv} (x), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right) \right) \right]$  for all  $m_1 m_2 \cdots m_r, n_1 n_2 \cdots n_s \in \mathbb{N}$  and scalars  $\langle \alpha_{m_1 m_2 \cdots m_r, n_1 n_2 \cdots n_s} \rangle$  with  $|\alpha_{m_1 m_2 \cdots m_r, n_1 n_2 \cdots n_s}| \leq 1$  for all  $m_1 m_2 \cdots m_r, n_1 n_2 \cdots n_s \in \mathbb{N}$ .

**Theorem 3.5.** *The spaces  $\left[ \chi_{mn}^{2Rf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  and  $\left[ \chi_{mn}^{2Bf} (q), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  are monotone.*

**Proof:**The proof follows from the Remark 2.6 and Theorem 3.4.

**Theorem 3.6.** *Let  $f_1$  and  $f_2$  be multiple sequence of Musielak modulus functions. Then we have*

$$(1) \left[ \chi_{mn}^{2Rf_1}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] \subseteq \left[ \chi_{mn}^{2Rf_2 \circ f_1}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$$

$$(2) \left[ \chi_{mn}^{2Rf_1}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] \cap \left[ \chi_{mn}^{2Rf_2}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] \subseteq$$

$$\left[ \chi_{mn}^{2Rf_1 + f_2}, \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$$

$$(3) \left[ \chi_{mn}^{2Rf_1} (q_1), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] \cap \left[ \chi_{mn}^{2Rf_1} (q_2), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] \subseteq$$

$\left[ \chi_{mn}^{2Rf_1} (q_1 + q_2), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right]$  where  $q_1$  and  $q_2$  are two semi norms.

(4) If  $q_1$  is stronger than  $q_2$ , then

$$\left[ \chi_{mn}^{2Rf_1} (q_1), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right] \subseteq \left[ \chi_{mn}^{2Rf_1} (q_2), \|(d(x_1), d(x_2), \dots, d(x_{n-1}))\|_p \right].$$

**Proof:** We have to take one condition and then easily prove to other condition. Hence it is trivial. Therefore omit the proof.

## COMPETING INTERESTS

The authors declare that there is no conflict of interests regarding the publication of this research paper.

## ACKNOWLEDGEMENT

The research work of the second author Deepmala is supported by the Science and Engineering Research Board (SERB), Government of India under SERB NPDF scheme, File Number: PDF/2015/000799. The third author wish to thank the Department of Science and Technology, Government of India for the financial sanction towards this work under FIST program SR/FST/MSI-107/2015.

## REFERENCES

- [1] T.J.F.A. Bromwich, An introduction to the theory of infinite series, Macmillan and Co.Ltd., New York, (1965).
- [2] G.H. Hardy, On the convergence of certain multiple series, Proc. Camb. Phil. Soc., 19 (1917), 86-95.
- [3] F. Moricz, Extentions of the spaces  $c$  and  $c_0$  from single to double sequences, Acta. Math. Hung., 57(1-2), (1991), 129-136.
- [4] F. Moricz and B.E. Rhoades, Almost convergence of double sequences and strong regularity of summability matrices, Math. Proc. Camb. Phil. Soc., 104, (1988), 283-294.
- [5] M. Basarir and O. Solanacan, On some double sequence spaces, J. Indian Acad. Math., 21(2) (1999), 193-200.
- [6] B.C. Tripathy, On statistically convergent double sequences, Tamkang J. Math., 34(3), (2003), 231-237.
- [7] A. Turkmenoglu, Matrix transformation between some classes of double sequences, J. Inst. Math. Comp. Sci. Math. Ser., 12(1), (1999), 23-31.
- [8] A. Gökhan and R. Çolak, The double sequence spaces  $c_2^P(p)$  and  $c_2^{PB}(p)$ , Appl. Math. Comput., 157(2), (2004), 491-501.
- [9] A. Gökhan and R. Çolak, Double sequence spaces  $\ell_2^\infty$ , ibid., 160(1), (2005), 147-153.
- [10] M. Zeltser, Investigation of Double Sequence Spaces by Soft and Hard Analytical Methods, Dissertationes Mathematicae Universitatis Tartuensis 25, Tartu University Press, Univ. of Tartu, Faculty of Mathematics and Computer Science, Tartu, 2001.
- [11] M. Mursaleen and O.H.H. Edely, Statistical convergence of double sequences, J. Math. Anal. Appl., 288(1), (2003), 223-231.
- [12] B. Altay and F. Başar, Some new spaces of double sequences, J. Math. Anal. Appl., 309(1), (2005), 70-90.
- [13] F. Başar and Y. Sever, The space  $\mathcal{L}_p$  of double sequences, Math. J. Okayama Univ, 51, (2009), 149-157.
- [14] N. Subramanian and U.K. Misra, The semi normed space defined by a double gai sequence of modulus function, Fasciculi Math., 46, (2010).
- [15] I.J. Maddox, Sequence spaces defined by a modulus, Math. Proc. Cambridge Philos. Soc., 100(1) (1986), 161-166.

- [16] J. Connor, On strong matrix summability with respect to a modulus and statistical convergence, *Canad. Math. Bull.*, 32(2), (1989), 194-198.
- [17] A. Pringsheim, Zurtheorie der zweifach unendlichen zahlenfolgen, *Math. Ann.*, 53, (1900), 289-321.
- [18] H.J. Hamilton, Transformations of multiple sequences, *Duke Math. J.*, 2, (1936), 29-60.
- [19] H.J. Hamilton, A Generalization of multiple sequences transformation, *Duke Math. J.*, 4, (1938), 343-358.
- [20] H.J. Hamilton, Preservation of partial Limits in Multiple sequence transformations, *Duke Math. J.*, 4, (1939), 293-297.
- [21] P.K. Kamthan and M. Gupta, Sequence spaces and series, Lecture notes, Pure and Applied Mathematics, 65 Marcel Dekker, Inc., New York , 1981.
- [22] J. Lindenstrauss and L. Tzafriri, On Orlicz sequence spaces, *Israel J. Math.*, 10 (1971), 379-390.
- [23] A. Wilansky, Summability through Functional Analysis, North-Holland Mathematical Studies, North-Holland Publishing, Amsterdam, Vol.85(1984).