

A short remark on the result of József Sándor

Yogesh J. Bagul¹

Department of Mathematics, K. K. M. College, Manwath,
Dist: Parbhani(M. S.) - 431505, India

ABSTRACT. We point out that Corollary 2.2 in the recently published article, 'On the Iyengar-Madhava Rao-Nanjundiah inequality and its hyperbolic version'[3] by József Sándor is slightly incorrect since its proof contains a gap. Fortunately, the proof can be corrected and this is the main aim of this note.

Keywords: Iyengar-Madhava Rao-Nanjundiah inequality.

2000 Mathematics subject classification: 26D05, 26D99; Secondary 26D07.

1. INTRODUCTORY REMARKS

The well known inequality [1, pp. 236], [2]

$$\cos x < \frac{\sin x}{x}; x \in (0, \pi/2) \quad (1.1)$$

has many applications in Mathematics. The inequality (1.1) has been studied and used extensively by many researchers in the literature. Its refinements and generalizations have been given by many others. Recently in [3] József Sándor proved the following statement:

Statement 1. ([3, Corollary 2.2]): The best constants c, d such that

$$\cos(x + c) < \frac{\sin x}{x} < \cos(x + d) \quad (1.2)$$

for $x \in (0, \pi/2)$ are $c = 0$ and $d = \arccos \frac{2}{\pi} - \frac{\pi}{2} \approx -0.690107$.

¹Corresponding author: yjbagul@gmail.com

Received: 06 February 2020

Accepted: 15 April 2020

The left inequality of (1.2) is undoubtedly valid; but right inequality in (1.2) is not valid in some portion of the interval $(0, \pi/2)$. This can be explained as follows:

In the proof of Statement 1, lastly József Sándor arrives at the conclusion that

$$x + d < \arccos\left(\frac{\sin x}{x}\right) < x \quad (1.3)$$

in $(0, \pi/2)$. Then he applies on (1.3) the cosine function considering it as decreasing in the intervals of values of functions in (1.3) which is logically incorrect for left inequality of (1.3) since $x + d$ is negative in the interval $(0, -d)$ whereas $\arccos(\sin x/x)$ is positive in the same interval. For applying cosine function on left inequality of (1.3), the values of the functions $x + d$ and $\arccos(\sin x/x)$ should lie in the same interval where cosine is increasing or decreasing. This general rule seems not to be considered by author of [3], so the result in Statement 1 is partially incorrect due to this simple mathematical mistake. However, it is worth noting that the technique of applying cosine function to obtain other inequalities in the same paper is interesting.

2. MAIN RESULT

We present a corrected version of Statement 1 in this section.

Theorem 2.1. *The best constants c, d such that*

$$\cos(x + c) < \frac{\sin x}{x}; \quad x \in (0, \pi/2) \quad (2.1)$$

and

$$\frac{\sin x}{x} < \cos(x + d); \quad x \in (\lambda, \pi/2) \quad (2.2)$$

where $\lambda \approx 0.43715$ are $c = 0$ and $d = \arccos \frac{2}{\pi} - \frac{\pi}{2} \approx -0.690107$.

Proof. The inequality (2.1) is already proved in [3]. So we prove only (2.2). As in the proof of [3, Corollary 2.2], consider the left inequality of (1.3) as

$$x + d < \arccos\left(\frac{\sin x}{x}\right)$$

where $d = \arccos \frac{2}{\pi} - \frac{\pi}{2} \approx -0.690107$. Clearly, in the interval $[-d, \pi/2)$ both the functions $x + d$ and $\arccos(\sin x/x)$ are non-negative. Therefore, applying strictly decreasing function cosine we get

$$\frac{\sin x}{x} < \cos(x + d); x \in [-d, \pi/2) \quad (2.3)$$

where $d \approx -0.690107$. Now it remains to consider the validity of (2.3) in $(0, -d)$. It is not difficult to check that $\cos(x + d)$ is strictly increasing and $\frac{\sin x}{x}$ is strictly decreasing in $(0, -d)$. Again the solution of equation $\cos(x + d) - \frac{\sin x}{x} = 0$ can be found by numerical methods techniques to be $x \approx 0.43715$. This shows that (2.3) is valid in $(\lambda, -d)$ also. The proof is complete. \square

The following graphical comparison of functions in (2.2) supports the proof.




FIGURE 1. Graphs of the functions in (2.2) for $x \in (0, \pi/2)$.

REFERENCES

- [1] D. S. Mitrinovi'c, *Analytic Inequalities*, Springer-Verlag, Berlin, 1970.
- [2] K. S. K. Iyengar, B. S. Madhava Rao and T. S. Nanjundiah, Some trigonometrical inequalities, *Half-Yearly J. Mysore Univ.*, **B(N.S.)(6)(1945)**, 1-12.
- [3] J. Sándor, On the Iyengar-Madhava Rao-Nanjundiah inequality and its hyperbolic version, *Notes on Number Theory and Discrete Mathematics*, **24(2)(2018)**, 134-139.