

## Solving fuzzy linear programming problems with linear membership functions-revisited

B. Farhadinia <sup>1</sup>

<sup>1</sup> Department of Mathematics, Quchan Institute of Engineering and Technology, Iran

**ABSTRACT.** Recently, Gasimov and Yenilmez proposed an approach for solving two kinds of fuzzy linear programming (FLP) problems. Through the approach, each FLP problem is first defuzzified into an equivalent crisp problem which is non-linear and even non-convex. Then, the crisp problem is solved by the use of the modified subgradient method. In this paper we will have another look at the earlier defuzzification process developed by Gasimov and Yenilmez in view of a perfectly acceptable remark in fuzzy contexts. Furthermore, it is shown that if the modified defuzzification process is used to solve FLP problems, some interesting results are appeared.

**Keywords:** Fuzzy linear programming problems; Modified subgradient method; Fuzzy decisive set method.

*2000 Mathematics subject classification:* xxxx, xxxx; Secondary xxxx.

### 1. INTRODUCTION

Since Gasimov and Yenilmez [6] investigated an approach for solving fuzzy linear programming (FLP) problems, it has been used frequently by a number of authors in various fields [5, 7]. Through Gasimov and Yenilmez's approach a FLP problem with fuzzy technological coefficients and fuzzy right-hand-side numbers corresponds to a crisp problem using the defuzzification process, known as the symmetric method of Bellman

---

<sup>1</sup> Corresponding author: bfarhadinia@yahoo.com.au  
Received: 26 September 2012  
Revised: 8 October 2013  
Accepted: 20 January 2014

and Zadeh [1]. Then, the modified subgradient method is applied for solving the obtained crisp problem.

Our main purpose of this paper is to present a revised formula for the membership function of fuzzy constraints with respect to a perfectly acceptable remark in fuzzy contexts. We will compare and show that this revision has some advantages rather than the Gasimov and Yenilmez's definition from points of view of (i) The number of required iterations to get the desired solution is reduced. (ii) The maximum satisfaction degree of the fuzzy decision set, that is, the objective of NLP problem,  $\lambda$ , reaches a more accurate optimum. (iii) The sequence  $\{\|g(x_k)\|\}$ , which evaluates how much constraints are violated, is controlled by a smaller upper bound.

The organization of this paper is as follows. Section 2 is devoted to recall a defuzzification process, known as the symmetric method of Bellman and Zadeh [1]. Also the proposed revision of defuzzification process is given in Section 2. The modified subgradient method and fuzzy decisive set method are presented algorithmically in Section 3. Finally, some comparative examples are provided in Section 4 to verify the main assertion of this contribution.

## 2. FLP PROBLEM AND DEFUZZIFICATION PROCESS

In this section, we restrict our attention to the following FLP problem involving fuzzy technological coefficients and fuzzy right-hand-side numbers.

$$\begin{aligned}
 \text{(FLP)} \quad & \max \quad \sum_{j=1}^n c_j x_j \\
 & \text{s.t.} \quad \sum_{j=1}^n \tilde{a}_{ij} x_j \leq \tilde{b}_i, \quad 1 \leq i \leq m, \\
 & \quad \quad \quad x_j \geq 0,
 \end{aligned}$$

where at least one  $x_j > 0$ , for  $j = 1, \dots, n$ .

In the sequel, we shall state that all fuzzy numbers in (FLP) are supposed to be described by linear membership functions. Hence, We assume that  $\tilde{a}_{ij}$  and  $\tilde{b}_i$  are fuzzy numbers with the following membership functions (for  $x \in \mathfrak{R}$ ):

$$\mu_{a_{ij}}(x) = \begin{cases} 1, & x < a_{ij}, \\ \frac{1}{d_{ij}}(a_{ij} + d_{ij} - x), & a_{ij} \leq x < a_{ij} + d_{ij}, \\ 0, & a_{ij} + d_{ij} \leq x, \end{cases} \quad (2.1)$$

$$\mu_{b_i}(x) = \begin{cases} 1, & x < b_i, \\ \frac{1}{p_i}(b_i + p_i - x), & b_i \leq x < b_i + p_i, \\ 0, & b_i + p_i \leq x. \end{cases} \quad (2.2)$$

In order to defuzzify (FLP), we can now proceed as follows. Firstly, we shall obtain the lower and upper bounds of the optimal values which are referred to as  $z_l$  and  $z_u$ , respectively.

Consider the four standard LP problems as follows:

$$(LP\_1) \quad z_1 = \max \sum_{j=1}^n c_j x_j \\ \text{s.t.} \quad \sum_{j=1}^n (a_{ij} + d_{ij}) x_j \leq b_i, \quad 1 \leq i \leq m, \\ x_j \geq 0,$$

$$(LP\_2) \quad z_2 = \max \sum_{j=1}^n c_j x_j \\ \text{s.t.} \quad \sum_{j=1}^n a_{ij} x_j \leq b_i + p_i, \quad 1 \leq i \leq m, \\ x_j \geq 0,$$

$$(LP\_3) \quad z_3 = \max \sum_{j=1}^n c_j x_j \\ \text{s.t.} \quad \sum_{j=1}^n (a_{ij} + d_{ij}) x_j \leq b_i + p_i, \quad 1 \leq i \leq m, \\ x_j \geq 0,$$

$$(LP\_4) \quad z_4 = \max \sum_{j=1}^n c_j x_j \\ \text{s.t.} \quad \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad 1 \leq i \leq m, \\ x_j \geq 0.$$

In the case that all the above LP problems have the finite optimal values, choosing the technological coefficient from the interval  $[a_{ij}, a_{ij} + d_{ij}]$  and the right-hand-side numbers from the interval  $[b_i, b_i + p_i]$  guarantees that the value of the objective function  $\sum_{j=1}^n c_j x_j$  is in the interval  $[z_l, z_u]$  where  $z_l = \min\{z_i, i = 1, 2, 3, 4\}$  and  $z_u = \max\{z_i, i = 1, 2, 3, 4\}$ .

Based on the above arguments, we may define the fuzzy set of optimal values  $G$  as follows:

$$\mu_G(x) = \begin{cases} 0, & \sum_{j=1}^n c_j x_j < z_l, \\ \frac{\sum_{j=1}^n c_j x_j - z_l}{z_u - z_l}, & z_l \leq \sum_{j=1}^n c_j x_j < z_u, \\ 1, & z_u \leq \sum_{j=1}^n c_j x_j. \end{cases} \quad (2.3)$$

Now we are in a position to give two different characterizations of the fuzzy set of the  $i$ -th constraint  $C_i$ . The first one is *the Gasimov and Yenilmez's definition (GD)* [3] and the second one is a new definition that we refer to it as *the revised definition (RD)*. The Gasimov and Yenilmez's definition for the fuzzy set of  $i$ -th constrain  $C_i$  is as

$$GD: \quad \mu_{C_i}(x) = \begin{cases} 0, & b_i < \sum_{j=1}^n a_{ij} x_j, \\ \frac{b_i - \sum_{j=1}^n a_{ij} x_j}{\sum_{j=1}^n d_{ij} x_j + p_i}, & \sum_{j=1}^n a_{ij} x_j \leq b_i < \sum_{j=1}^n (a_{ij} + d_{ij}) x_j + p_i, \\ 1, & \sum_{j=1}^n (a_{ij} + d_{ij}) x_j + p_i \leq b_i. \end{cases}$$

Before attempting to present the revised definition (RD), we recall a perfectly acceptable remark in fuzzy contexts. (See [1, 10]).

**Remark 1.** A proper fuzzy membership function should be set 0 if the constraints are strongly violated in the crisp sense, and 1 if they are very well satisfied in the crisp sense. Also it should increase monotonously from 0 to 1.

As can be seen, (LP\_1), (LP\_2), (LP\_3) and (LP\_4) have the same objective function, but their constraints are different. By shifting the term  $P = (p_i)_{1 \times m} \geq 0$  in all constraints to the left-hand side, one gets

$$\begin{aligned}(A + D)x &\leq b, \\ Ax - P &\leq b, \\ (A + D)x - P &\leq b, \\ Ax &\leq b,\end{aligned}$$

where  $x = (x_i)_{n \times 1}$ ,  $b = (b_i)_{1 \times m}$ ,  $A = (a_{ij})_{m \times n}$  and  $D = (d_{ij})_{m \times n} \geq 0$ . For any  $x \geq 0$ , we define

$$\overline{b(x)} = \max\{(A + D)x, Ax - P, (A + D)x - P, Ax\}, \quad (2.4)$$

$$\underline{b(x)} = \min\{(A + D)x, Ax - P, (A + D)x - P, Ax\}. \quad (2.5)$$

Hence, by virtue of Remark 1 and (2.4)-(2.5), a fuzzy constraint  $C_i$  should be characterized by

$$\mu_{C_i}(x) = \begin{cases} 0, & b < \underline{b(x)}, \\ \in [0, 1], & \underline{b(x)} \leq b \leq \overline{b(x)}, \\ 1, & \overline{b(x)} \leq b. \end{cases} \quad (2.6)$$

One can easily verify that for any  $x \geq 0$ ,

$$\overline{b(x)} = (A + D)x, \quad (2.7)$$

$$\underline{b(x)} = Ax - P. \quad (2.8)$$

Consequently, by virtue of  $\mu_{C_i}(x)$  defined in (2.6) and (2.7)-(2.8), the revised definition for a fuzzy constraint,  $C_i$ , may be defined by

$$\text{RD : } \mu_{C_i}(x) = \begin{cases} 0, & b_i < \sum_{j=1}^n a_{ij}x_j - p_i, \\ \frac{b_i - \sum_{j=1}^n a_{ij}x_j + p_i}{\sum_{j=1}^n d_{ij}x_j + p_i}, & \sum_{j=1}^n a_{ij}x_j - p_i \leq b_i < \sum_{j=1}^n (a_{ij} + d_{ij})x_j, \\ 1, & \sum_{j=1}^n (a_{ij} + d_{ij})x_j \leq b_i. \end{cases}$$

Now, by making use of the definition of the fuzzy decision proposed by Bellman and Zadeh [1], we can characterize the fuzzy decision set  $D$  as follows:

$$D = G \cap \left\{ \bigcap_{i=1}^m C_i \right\}.$$

In this regards, an optimum solution can be selected as the design for which one may get the maximum of the membership function. That is,

$$\mu_D(x^*) = \max_{x \geq 0} \{\mu_D(x)\}, \quad (2.9)$$

where  $\mu_D(x) = \min_{x \geq 0} \{\mu_G(x), \mu_{C_1}(x), \dots, \mu_{C_m}(x)\}$ . (See [1, 10]).

Suppose that  $\lambda = \mu_D(x)$ . Therefore, the optimization problem (2.9) can be restated in the form of

$$\begin{aligned} \text{(D-SET)} \quad & \max \quad \lambda \\ & \text{s.t.} \quad \mu_G(x) \geq \lambda, \\ & \quad \mu_{C_i}(x) \geq \lambda, \quad 1 \leq i \leq m, \\ & \quad x \geq 0, \\ & \quad 0 \leq \lambda \leq 1. \end{aligned}$$

Putting together the definition of objective function  $G$  in (2.3), the two different definitions of constraints  $C_i$  labeled by GD and RD and the optimization problem (D-SET), one can defuzzify (FLP) into the two following non-convex optimization problems:

$$\begin{aligned} \text{(P\_GD)} \quad & \max \quad \lambda \\ & \text{s.t.} \quad \lambda(z_u - z_l) - \sum_{j=1}^n c_j x_j + z_l \leq 0, \\ & \quad \sum_{j=1}^n (a_{ij} + \lambda d_{ij}) x_j + \lambda p_i - b_i \leq 0, \quad 1 \leq i \leq m, \\ & \quad x \geq 0, \\ & \quad 0 \leq \lambda \leq 1, \end{aligned}$$

and

$$\begin{aligned} \text{(P\_RD)} \quad & \max \quad \lambda \\ & \text{s.t.} \quad \lambda(z_u - z_l) - \sum_{j=1}^n c_j x_j + z_l \leq 0, \\ & \quad \sum_{j=1}^n (a_{ij} + \lambda d_{ij}) x_j + \lambda p_i - p_i - b_i \leq 0, \quad 1 \leq i \leq m, \\ & \quad x \geq 0, \\ & \quad 0 \leq \lambda \leq 1, \end{aligned}$$

Needless to say that the non-convexity of (P\_GD) and (P\_RD) is due to the presence of  $\lambda x_j$  in their constraints.

### 3. TWO METHODS FOR SOLVING NON-CONVEX OPTIMIZATION PROBLEMS

Let us emphasize that this section will not contain all definitions and theorems which are required for implementing both the well-known methods: the modified subgradient method [2] and the fuzzy decisive set method [8]. The interested reader is referred to [2, 3] and [8].

In the modified subgradient algorithm instead of focussing on the primal problem, it focuses on the dual problem obtained with respect to the sharp Lagrangian. Let us consider the following primal mathematical programming problem:

$$(P\_I) \quad \text{minimize } f(x) \text{ over all } x \in S \text{ satisfying } g(x) = 0,$$

where  $S$  is a compact subset of  $\mathfrak{R}^n$ , and both functions  $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$  and  $g : \mathfrak{R}^n \rightarrow \mathfrak{R}^m$  are continuous. Let  $\mathfrak{R}^+$ ,  $\|\cdot\|$  and  $\langle \cdot, \cdot \rangle$  be the set of nonnegative real numbers, the Euclidean norm and the Euclidean inner product on  $\mathfrak{R}^m$ , respectively. The augmented Lagrangian  $L : \mathfrak{R}^n \times \mathfrak{R}^m \times \mathfrak{R}^+ \rightarrow \mathfrak{R}$  associated with (P\_I) is defined in the form of

$$L(x, u, c) = f(x) + c\|g(x)\| - \langle u, g(x) \rangle, \quad (3.1)$$

where  $x \in \mathfrak{R}^n$ ,  $u \in \mathfrak{R}^m$  and  $c \in \mathfrak{R}^+$ .

The dual function  $H : \mathfrak{R}^m \times \mathfrak{R}^+ \rightarrow \mathfrak{R}$  is defined in the form of

$$H(u, c) = \min_{x \in S} \{f(x) + c\|g(x)\| - \langle u, g(x) \rangle\}. \quad (3.2)$$

By virtue of (11), we define the dual problem of (P\_I) by

$$(P\_II) \quad \text{minimize } H(u, c) \text{ over all } (u, c) \in \mathfrak{R}^m \times \mathfrak{R}^+.$$

Theorems 1,2,3 and 4 in [3] show that zero duality gap and saddle point properties hold. Theorem 5 in [3] which is restated below is used to define a stopping criteria for the modified subgradient algorithm.

**Theorem 1.** Suppose that (P\_I) and (P\_II) have a finite solution and assume that for some  $(\bar{u}, \bar{c}) \in \mathfrak{R}^m \times \mathfrak{R}^+$ , and  $\bar{x} \in S$ ,

$$\min_{x \in S} L(x, \bar{u}, \bar{c}) = f(\bar{x}) + \bar{c}\|g(\bar{x})\| - \langle \bar{u}, g(\bar{x}) \rangle.$$

Then,  $\bar{x}$  is a solution to (P\_I) and  $(\bar{u}, \bar{c})$  is a solution to (P\_II) if and only if

$$g(\bar{x}) = 0.$$

We now outline the modified subgradient algorithm as follows.

### Modified Subgradient Algorithm

**Initialization Step.** Choose  $(u^0, c^0)$  with  $c^0 \geq 0$ . Set  $k = 0$ .

**Main Step.** Given  $(u^k, c^k)$ :

**Step 1.** Solve the following subproblem:

$$\min_{x \in S} \{f(x) + c^k\|g(x)\| - \langle u^k, g(x) \rangle\}.$$

Let  $x^k$  be a solution. If  $g(x^k) = 0$ , then stop and by Theorem 1

$(u^k, c^k)$  is a solution of (P\_II) and  $x^k$  is solution of (P\_I).

**Step 2.** Set

$$u^{k+1} = u^k - s^k g(x^k),$$

$$c^{k+1} = c^k + (s^k + \epsilon^k) \|g(x^k)\|,$$

where  $s^k, \epsilon^k > 0$ . Set  $k = k + 1$  and repeat Main Step.

The convergence results of the modified subgradient algorithm are given in [6].

Here, we outline another algorithm known as the fuzzy decisive set method [8] and is implemented for solving problems in the form of

$$\max\{\lambda \mid G(x, \lambda) = 0 \text{ and } x \geq 0, 0 \leq \lambda \leq 1\}.$$

#### Fuzzy Decisive Set Algorithm

**Initialization Step.** Set  $k = 0$ . Let  $\lambda_0 = 1$ . If there exists a set which satisfies  $G(x, \lambda_0) = 0$  and  $x \geq 0$  then,  $\lambda^* = \lambda_0$ . If this is not the case, set  $\lambda_0^L = 0$  and  $\lambda_0^R = 1$  and go to Main Step.

**Main Step.** Set  $k = k + 1$ , and let  $\lambda_k = \frac{\lambda_{k-1}^L + \lambda_{k-1}^R}{2}$ .

If  $G(x, \lambda_k) = 0$  does not hold for a  $x \geq 0$ , then set  $\lambda_k^L = \lambda_k$  and  $\lambda_k^R = \lambda_{k-1}^R$ .

If  $G(x, \lambda_k) = 0$  does hold for a  $x \geq 0$ , then set  $\lambda_k^R = \lambda_k$  and  $\lambda_k^L = \lambda_{k-1}^L$ .

Set  $k = k + 1$ . If  $|\lambda_{k+1} - \lambda_k| < \epsilon$ , where  $\epsilon > 0$  is a small constant, then stop and output  $\lambda^* = \lambda_{k+1}$ .

#### 4. COMPARISON RESULTS

In order to apply the modified subgradient algorithm for solving (P\_GD) and (P\_RD), they should be transformed into the (P\_I) form. Hence, by the use of slack variables  $q_i^r$ , for  $r = 1, 2$ ,  $i = 0, 1, \dots, m$ , and the relation  $\max \lambda = -\min(-\lambda)$ , problems (P\_GD) and (P\_RD) may be

restated as

$$\begin{aligned}
 (\text{P\_GD}) \quad & \max \quad \lambda = -\min(-\lambda) \\
 & \text{s.t.} \quad g_0^1(x, \lambda, q_0^1) = \lambda(z_u - z_l) - \sum_{j=1}^n c_j x_j + z_l + q_0^1 = 0, \\
 & \quad g_i^1(x, \lambda, q_i^1) = \sum_{j=1}^n (a_{ij} + \lambda d_{ij}) x_j + \lambda p_i - b_i + q_i^1 = 0, \\
 & \quad 1 \leq i \leq m, \\
 & \quad x \geq 0, \quad q_0^1, q_i^1 \geq 0, \quad 0 \leq \lambda \leq 1,
 \end{aligned}$$

and

$$\begin{aligned}
 (\text{P\_RD}) \quad & \max \quad \lambda = -\min(-\lambda) \\
 & \text{s.t.} \quad g_0^2(x, \lambda, q_0^2) = \lambda(z_u - z_l) - \sum_{j=1}^n c_j x_j + z_l + q_0^2 = 0, \\
 & \quad g_i^2(x, \lambda, q_i^2) = \sum_{j=1}^n (a_{ij} + \lambda d_{ij}) x_j + \lambda p_i - p_i - b_i + q_i^2 = 0, \\
 & \quad 1 \leq i \leq m, \\
 & \quad x \geq 0, \quad q_0^2, q_i^2 \geq 0, \quad 0 \leq \lambda \leq 1,
 \end{aligned}$$

where

$$S^1 = \{(x, \lambda, q^1) \mid x = (x_1, \dots, x_n), q^1 = (q_0^1, q_1^1, \dots, q_m^1), x_j \geq 0, q_i^1 \geq 0, 0 \leq \lambda \leq 1\},$$

$$S^2 = \{(x, \lambda, q^2) \mid x = (x_1, \dots, x_n), q^2 = (q_0^2, q_1^2, \dots, q_m^2), x_j \geq 0, q_i^2 \geq 0, 0 \leq \lambda \leq 1\}.$$

In what follows, we will apply the modified subgradient algorithm in accordance with the Gasimov and Yenilmez's definition **GD** and the revised definition **RD** for a fuzzy constraint to some test problems.

For the comparison of the results obtained by the use of the Gasimov and Yenilmez's definition **GD** and the revised definition **RD**, we solve the (FLP) given in [3] with respect to the cases in which  $(p_i) = [p_1, p_2]^\top$  are, but may not be limited to,

$$\text{Case 1. } (p_i) = [2.9, 3.9]^\top,$$

$$\text{Case 2. } (p_i) = [8, 10]^\top.$$

Now, consider the following (FLP) discussed in [3]

$$\begin{aligned}
 \max \quad & x_1 + x_2 \\
 \text{s.t.} \quad & \tilde{1}x_1 + \tilde{2}x_2 \leq \tilde{3} \\
 & \tilde{2}x_1 + \tilde{3}x_2 \leq \tilde{4} \\
 & x_1, x_2 \geq 0,
 \end{aligned}$$



where fuzzy parameters  $\tilde{1} = L(1, 1)$ ,  $\tilde{2} = L(2, 1)$ ,  $\tilde{3} = L(3, 2)$ ,  $\tilde{b}_1 = \tilde{3} = L(3, p_1)$  and  $\tilde{b}_2 = \tilde{4} = L(4, p_2)$  are taken as defined in [9]. That is,

$$(a_{ij}) = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}, \quad (d_{ij}) = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \quad \Longrightarrow \quad (a_{ij} + d_{ij}) = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix},$$

$$(b_i) = \begin{bmatrix} 3 \\ 4 \end{bmatrix}, \quad (p_i) = \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} \quad \Longrightarrow \quad (b_i + p_i) = \begin{bmatrix} 3 + p_1 \\ 4 + p_2 \end{bmatrix}.$$

For more convenience, we use the following notations to interpret the results which have been reported in Tables 1-4.

- $k$  is the number of iteration,
- $(u^k, c^k)$  is a vector of Lagrange multipliers at  $k$ -th iteration,
- $x^k$  is a minimizer of Lagrange function  $L(x, u^k, c^k)$  over  $x \in S^r$ ,  $r = 1, 2$ ,
- $\bar{H}$  is the upper bound for the values of dual function,
- $s^k$  is the stepsize parameter calculated at the  $k$ -th iteration by the formula  $s^k = \frac{\bar{H} - H(u^k, c^k)}{5\|g(x^k)\|^2}$ ,
- $\epsilon^k = 0.95 s^k$ .

We have taken  $\|g(x^k)\| \leq 10^{-5}$ , as the stopping criteria in each example.

**Example 1.** (*Case 1.*) Let

$$(p_i) = \begin{bmatrix} 2.9 \\ 3.9 \end{bmatrix} \quad \Longrightarrow \quad (b_i + p_i) = \begin{bmatrix} 5.9 \\ 7.9 \end{bmatrix}.$$

For solving (FLP) in this case, we must solve the two subproblems (LP\_1) and (LP\_2) which are expressed by the use of (2.7) and (2.8) as follows:

$$\begin{array}{ll} z_1 = \max & x_1 + x_2 \\ \text{s.t.} & 2x_1 + 3x_2 \leq 3, \\ & 4x_1 + 5x_2 \leq 4, \\ & x_1, x_2 \geq 0, \end{array} \qquad \begin{array}{ll} z_2 = \max & x_1 + x_2 \\ \text{s.t.} & x_1 + 2x_2 \leq 5.9, \\ & 2x_1 + 3x_2 \leq 7.9, \\ & x_1, x_2 \geq 0. \end{array}$$

The optimal solutions are

$$\begin{array}{ll} z_1^* = 1, & z_2^* = 3.95, \\ (x_1^*, x_2^*) = (1, 0), & (x_1^*, x_2^*) = (3.95, 0). \end{array}$$

Now, bearing the solution of the latter subproblems in mind, we establish problems ((P\_GD)) and ((P\_RD)) in the forms of

$$\begin{aligned}
 ((P\_GD)) \quad & \max \quad \lambda \\
 & s.t. \quad -x_1 - x_2 + 2.95\lambda + 1 + q_0^1 = 0, \\
 & \quad (1 + \lambda)x_1 + (2 + \lambda)x_2 + 2.9\lambda - 3 + q_1^1 = 0, \\
 & \quad (2 + 2\lambda)x_1 + (3 + 2\lambda)x_2 + 3.9\lambda - 4 + q_2^1 = 0, \\
 & \quad 1 \leq x_1 \leq 3.95, \quad 0 \leq x_2 \leq 0, \quad 0 \leq \lambda \leq 1,
 \end{aligned}$$

and

$$\begin{aligned}
 ((P\_RD)) \quad & \max \quad \lambda \\
 & s.t. \quad -x_1 - x_2 + 2.95\lambda + 1 + q_0^2 = 0, \\
 & \quad (1 + \lambda)x_1 + (2 + \lambda)x_2 + 2.9\lambda - 5.9 + q_1^2 = 0, \\
 & \quad (2 + 2\lambda)x_1 + (3 + 2\lambda)x_2 + 3.9\lambda - 7.9 + q_2^2 = 0, \\
 & \quad 1 \leq x_1 \leq 3.95, \quad 0 \leq x_2 \leq 0, \quad 0 \leq \lambda \leq 1.
 \end{aligned}$$

We have solved ((P\_GD)) and ((P\_RD)) firstly by using the fuzzy decisive set method and secondly by using the modified subgradient method.

- By the use of the fuzzy decisive set method, the solution of both problems ((P\_GD)) and ((P\_RD)) are obtained at the twenty first iterations, but with different optimal values  $\lambda_{20}((P\_GD)) = 0.1574$  and  $\lambda_{20}((P\_RD)) = 0.4142$ .
- The results obtained by the use of the modified subgradient method are illustrated in Table 1 and Table 2.

Table 1. The results of using the modified subgradient method for solving ((P\_GD))

k	$u_0^k$	$u_1^k$	$u_2^k$	$c^k$	$x_1^k$	$x_2^k$	$\lambda$	$\bar{H}$	$H_k$	$\ g(x_k)\ $	$s^k$
1	0	0	0	0	1	0	1	0	-1	5.2462	0.0073
2	-0.0214	-0.0138	-0.0283	0.0743	1	0	1	0	-0.4100	5.2462	0.0029
3	-0.0302	-0.0195	-0.0400	0.1048	1	0	0.5102	0	-0.2345	1.8129	0.0142
4	-0.0517	-0.0193	-0.0544	0.1553	1.4635	0	0.1571	0	-	$5.2 \times 10^{-6}$	-

Table 2. The results of using the modified subgradient method for solving ((P\_RD))

k	$u_0^k$	$u_1^k$	$u_2^k$	$c^k$	$x_1^k$	$x_2^k$	$\lambda$	$\bar{H}$	$H_k$	$\ g(x_k)\ $	$s^k$
1	0	0	0	0	1	0	1	0	-1	3.1149	0.0206
2	-0.0608	-0.0206	0	0.1252	0.1257	0	1	0	-0.4800	2.9224	0.0112
3	-0.0910	0.0152	-0.1159	0.1893	2.2219	0	0.4142	0	-	$1.3 \times 10^{-6}$	-

Once again remember that problems ((P\_GD)) and ((P\_RD)) have been generated according to the Gasimov and Yenilmez's definition GD and the revised definition RD for a fuzzy constraint, respectively.

**Discussion.** By comparing the results reported in Table 1 and Table 2, the following observations are evident: (i) The number of iterations for solving ((P\_RD)),  $k = 3$ , is less than one for solving ((P\_GD)),  $k = 4$ . (ii) The satisfaction level of constraints in the problem ((P\_RD)) is more desirable than the counterpart in ((P\_GD)), because  $\|g(x_k)\| = 1.3 \times 10^{-6}$  for ((P\_RD)) is less than  $\|g(x_k)\| = 5.2 \times 10^{-6}$  for ((P\_GD)). (iii) The maximum satisfaction degree of fuzzy decision set, that is, the optimum of ((P\_RD)),  $\lambda = 0.4142$ , has been more improved rather than the counterpart of ((P\_GD)),  $\lambda = 0.1571$ .

We note that all the above observations are in agreement with the results of the next experiment.

By the same manner as described in Example 1, we have examined the next example and consequently the results are summarized in Table 3 and Table 4.

**Example 2.** (*Case 2.*) Let

$$(p_i) = \begin{bmatrix} 8 \\ 10 \end{bmatrix} \implies (b_i + p_i) = \begin{bmatrix} 11 \\ 14 \end{bmatrix}.$$

We have solved ((P\_GD)) and ((P\_RD)) firstly by using the fuzzy decisive set method and secondly by using the modified subgradient method.

- By the use of the fuzzy decisive set method, the solution of problems ((P\_GD)) and ((P\_RD)) are obtained at the twenty four first and at the sixteen first iterations, respectively, while optimal values are  $\lambda_{24}((P_GD)) = 0.0801$  and  $\lambda_{16}((P_RD)) = 0.4143$ .
- The results obtained by the use of the modified subgradient method are illustrated in Table 3 and Table 4.

Table 3. The results of using the modified subgradient method for solving ((P\_GD))

k	$u_0^k$	$u_1^k$	$u_2^k$	$c^k$	$x_1^k$	$x_2^k$	$\lambda$	$\bar{H}$	$H_k$	$\ g(x_k)\ $	$s^k$
1	0	0	0	0	1	0	1	0	-1	13.6015	0.0011
2	-0.0065	0.0076	-0.0108	0.0287	1	0	1	0	-0.4100	13.6015	0.0004
3	-0.0091	-0.0107	-0.0152	0.0404	1	0	1	0	-0.1681	13.6015	0.0002
4	-0.0102	-0.0119	-0.0176	0.0452	1	0	0.2755	0	-0.1328	2.1600	0.0057
5	-0.0196	-0.0147	-0.0245	0.0692	1	0	0.1900	0	-0.8131	1.2100	0.0111
6	-0.0323	-0.0115	-0.0276	0.0954	1.4807	0	0.0801	0	-	$1.4 \times 10^{-6}$	-

Table 4. The results of using the modified subgradient method for solving ((P\_RD))

k	$u_0^k$	$u_1^k$	$u_2^k$	$c^k$	$x_1^k$	$x_2^k$	$\lambda$	$\bar{H}$	$H_k$	$\ g(x_k)\ $	$s^k$
1	0	0	0	0	1	0	1	0	-1	6.0828	0.0054
2	-0.0324	0.0054	0	0.0641	8.4852	0	0.4143	0	-0.4420	$0.2 \times 10^{-6}$	-

## 5. CONCLUSIONS

In this article we have suggested a revised formula for the membership function of fuzzy constraints involved in fuzzy linear programming problems with fuzzy technological coefficients and fuzzy right-hand side. Comparing the results obtained based on the Gasimov and Yenilmez's formula and the revised formula indicates that the proposed formula has some advantages such as: the number of required iterations to get the desired solution is reduced; the maximum satisfaction degree of the fuzzy decision set reaches a more accurate optimum; and the sequence which evaluates how much constraints are violated, is controlled by a smaller upper bound.

## REFERENCES

- [1] R.E. Bellman & L.A. Zadeh. "Decision making in a fuzzy environment". *Management Science*. Volume 17, 141-164. (1970).
- [2] R.N. Gasimov & A.M. Rubinov. "On augmented Lagrangians for optimization problems with a single constraint". *J. Global Optimization*. Volume 28, 153-173. (2004).
- [3] R.N. Gasimov & K. Yenilmez. "Solving fuzzy linear programming problems with linear membership functions". *Turk. J. Math.*, Volume 26, 375-396. (2002).
- [4] G.J. Klir & B. Yuan. "Fuzzy sets and fuzzy logic-theory and applications". *Prentice-Hall Inc.*, (1995).
- [5] S. Mukherjee & Z. Chen & A. Gangopadhyay. "A fuzzy programming approach for data reduction and privacy in distance-based mining". *Int. J. Information and computer Security*, Volume 20, 27-47. (2008).
- [6] N. Rafail & R.N. Gasimov. "Augmented Lagrangian duality and nondifferentiable optimization methods in nonconvex programming". *J. Global Optimization*, Volume 24, 187-203. (2002).
- [7] N. Safaei & M.S. Mehrobad & R.T. Moghaddam & F. Sassani. "A fuzzy programming approach for a cell formation problem with dynamic and uncertain conditions". *Fuzzy Sets and Systems*, Volume 159, 215-236. (2008).
- [8] M. Sakawa & H. Yana. "Interactive decision making for multiobjective linear fractional programming problems with fuzzy parameters". *Cybernetics Systems*, Volume 16, 377-397. (1985).
- [9] T. Shaocheny. "Interval number and fuzzy number linear programming". *Fuzzy Sets and Systems*, Volume 66, 301-306. (1994).
- [10] H.J. Zimmermann. "Description and optimization of fuzzy systems". *Int. J. General systems*, Volume 2, 209-215. (1976).