

Application of The Sine-Gordon Expansion Method on Nonlinear Various Physical Models

Sait San, Bahri Koç and Sukri Khareng ¹

Department of Mathematics - Computer, 26480, Eskişehir - TURKEY

ABSTRACT. In this paper, by utilizing the Sine-Gordon expansion method, soliton solutions of the higher-order improved Boussinesq equation, Kuramoto-Sivashinsky equation, and seventh-order Sawada-Kotera equation are obtained. Given partial differential equations are reduced to ordinary differential equations, by choosing the compatible wave transformation associated with the structure of the equation. Based on the solution of the Sine-Gordon equation, a polynomial system of equations is obtained according to the principle of homogeneous balancing. The solution of the outgoing system gives the parameters which are included by the solution. Plot3d and Plot2d graphics are given in detail. As a result, many different graphic models are obtained from soliton solutions of equations that play a very important role in mathematical physics and engineering.

Keywords: The Sine-Gordon Expansion Method, Travelling Wave Solution, Nonlinear Equations, Higher Order Boussinesq equation.

¹Corresponding author: Email: ssan@ogu.edu.tr


Received: 02 September 2021

Revised: 23 September 2021

Accepted: 01 October 2021

How to Cite: San, Sait; Koc, Bahri; Khareng, Sukri. Application of The Sine-Gordon Expansion Method on Nonlinear Various Physical Models, *Casp.J. Math. Sci.*, **12**(1)(2023), 30-50.

This work is licensed under a Creative Commons Attribution 4.0 International License.

 Copyright © 2023 by University of Mazandaran. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

1. INTRODUCTION

Nonlinear partial differential equations are encountered in modeling of many problems in physics and engineering. In recent studies, the integrability, the existence and uniqueness of their solutions, obtaining conservation laws, finding numerical and exact solutions of these equations have been discussed. Since there is no universal method for obtaining exact solutions, it has caused many researchers to work in this field. Using the functionality and ease of symbolic computing programs such as maple matlab, many powerful techniques developed such as improved (G'/G) - expansion method [1], $\exp(-\Omega(\xi))$ -expansion function method [2], the generalized Riccati equation mapping method [3], modified trial equation method [4], homotopy perturbation method [5], Kudryashov method [6]- [9], Jacobi elliptic function method [10], lie symmetry analysis method [11]-[13] and so on. Most methods have been used to find different perspectives and to obtain different solutions. All these exact solution methods are effective and lead to assorted types of solutions such as trigonometric solutions, non-periodic solutions, soliton and solitary solutions. Especially the last two types are very important in nonlinear physical phenomena and engineering.

In this study, we considered the sine-gordan expansion method to find exact solutions of various physical models. Firstly, we examined the higher-order improved Boussinesq equation which were derived formally from 2D water wave problem. Schneider and Wayne introduced a class of Boussinesq equations to model the water wave problem with surface tension [14].

$$-u_{xxxxtt} + u_{xxtt} - u_{tt} + u_{xx} + \mu u_{xxxx} + (u)_{xx}^2 = 0 \quad (1.1)$$

They have showed that the equation (1.1) really can be expressed by two decoupled Kawahara equations for a degenerate case. In [15], Akçağıl and Gözükızıl utilized the tanh-coth method and obtained some travelling wave solutions. The authors of [16] have also showed that the finite time blow up solutions by employing the improved convexity method. And in [17], Wang and Xue considered the Cauchy problem for sixth order Boussinesq equation and proved that the existence and uniqueness of the local solution by the contraction mapping theorem.

Another physical model is Kuramoto-Sivashinsky (KS) equation is presented [18] as

$$u_t + auu_x + bu_{xx} + ku_{xxx} = 0. \quad (1.2)$$

in which a, b and k are some arbitrary constants. It appears in many scientific areas such as phase turbulence systems [19], an example of spatio temporal chaos [20], flame front propagation [21] and so on. In

[22], approximate analytical solutions were found with homotopy analysis method and compared numerical solutions. Khater et al. [23], Chebyshev spectral collocation method were performed and obtained approximate numerical solutions.

As the last application, the seventh-order Sawada-Kotera equation was examined [24].

$$u_t + [63u^4 + 63(2u^2u_{xx} + u(u_x)^2) + 21(uu_{xxxx} + (u_{xx})^2 + u_xu_{xxx}) + u_{xxxxxx}]_x = 0 \quad (1.3)$$

Higher order KdV type equations have many applications in fluid dynamics [25]-[29]. Jafari et al.[30] obtained approximate solutions by applying the Adomian decomposition method and He's variational iteration method for two different physical models of higher order KdV type equations. Wazwaz employed the tanh-coth method and Hirota's direct method and obtained multiple soliton solutions [31]. The authors of [32], performed Bell polynomial approach to investigate integrable properties and constructed Lax pair, Backlund Transformation, Hirota D-operators and infinite conservation laws.

2. METHODOLOGY OF THE METHOD

Firstly, we will give some preliminaries about the methodology of the Sine-Gordon expansion method [33]- [37]. This method is constructed on the well known Sine-Gordan equation.

$$u_{xx} - u_{tt} = m^2 \sin(u) \quad (2.1)$$

in which $u = u(x, t)$ is definition as a arbitrary function and m is a constant. By taking $u(x, t) = U(\xi)$ where it is applied $\xi = kx + lt$ which is called wave transformation, (2.1) equation is reduced in one dimension form.

$$U'' = \frac{m^2}{k^2 - l^2} \sin^2\left(\frac{U}{2}\right) \quad (2.2)$$

When multiply by U' and integrating once, we get a simpler equation.

$$\left[\left(\frac{U}{2}\right)'\right]^2 = \frac{m^2}{k^2 - l^2} \sin^2\left(\frac{U}{2}\right) + K \quad (2.3)$$

Here K is an integration constant. If we get $K = 0$, $\frac{U}{2} = w(\xi)$ and $\frac{m^2}{k^2 - l^2} = \alpha^2$, we get a simpler equation as follows.

$$w' = \alpha \sin(w) \quad (2.4)$$

Solution of Eq. (2.4) with taking $\alpha = 1$ can be find as below

$$\sin(w) = \sin(w(\xi)) = \frac{2pe^\xi}{p^2e^{2\xi} + 1} \Big|_{p=1} = \operatorname{sech}(\xi) \quad (2.5)$$

$$\cos(w) = \cos(w(\xi)) = \frac{2pe^\xi - 1}{p^2e^{2\xi} + 1} \Big|_{p=1} = \tanh(\xi), \quad (2.6)$$

where $p \neq 0$ is the integration constant.

We suppose that a nonlinear differential equation of the polynomial form

$$H(u, u_x, u_t, u_{xx}, u_{xt}, u_{tt}, \dots) = 0. \quad (2.7)$$

Applying the compatible wave transformation, (3.2) can be converted to ODE

$$O(U, U', U'', U''' \dots) = 0 \quad (2.8)$$

Thence, the expected solution to (3.3) of the form

$$U(\xi) = \sum_{i=1}^n \tanh^{i-1}(\xi) [B_i \operatorname{sech}(\xi) + A_i \tanh(\xi)] + A_0 \quad (2.9)$$

Thanks to (2.5) and (3.1) equations, we get equation (3.4) as following

$$U(\omega) = \sum_{i=1}^n \cos^{i-1}(\omega) [B_i \sin(\omega) + A_i \cos(\omega)] + A_0 \quad (2.10)$$

Operate the homogeneous balance principle to obtain the value of n . In equation (2.10), necessary derivatives are taken and replaced in equation (3.3). In obtained expression, letting the coefficients of $\sin^p \cos^q$ of equal power to be all zero, we construct an algebraic system. When the system is solved by Maple, the values of unknowns A_i, B_i, k, l are found. Finally, one can easily construct the soliton solutions of Eq. (3.3).

3. APPLICATIONS

3.1. Higher Order Boussinesq Equation. In this section, we start with the sixth-order Boussinesq equation,

$$-u_{xxxxxtt} + u_{xxtt} - u_{tt} + u_{xx} + \mu u_{xxxx} + u_{xx}^2 = 0, \quad (3.1)$$

where $u(x, t)$ is an analytic function and μ is arbitrary constant.

If we use the wave transformation $\xi = x - ct$ into Eq.(3.1), it transformed into an ODE

$$-c^2 u^{(vi)} + c^2 u'''' - c^2 u'' + u'' + \mu u'''' + 2(u')^2 + 2uu'' = 0, \quad (3.2)$$

where prime expresses differentiation with respect to ξ . If we balance between the highest order derivative term $u^{(vi)}$ and nonlinear term of the highest degree uu'' in Eq.(3.2), we obtain $n = 4$.

Thus, the exact solution of the related equation has the form,

$$\begin{aligned} u(\xi) = & A_0 + B_1 \sec h(\xi) + B_2 \sec h(\xi) \tanh(\xi) \\ & + B_3 \sec h(\xi) \tanh^2(\xi) + B_4 \sec h(\xi) \tanh^3(\xi) \\ & + A_1 \tanh(\xi) + A_2 \tanh^2(\xi) + A_3 \tanh^3(\xi) + A_4 \tanh^4(\xi), \end{aligned} \quad (3.3)$$

where $u(x, t) = U(\xi)$ satisfies Eq.(2.5) and $A_0, A_1, A_2, A_3, A_4, B_1, B_2, B_3, B_4$ are undetermined constants.

Firstly, we substitute Eq.(3.3) into Eq.(3.2) with taking the necessary derivatives and employ Eq.(2.5), in the end we equate all coefficients of the functions $[\cos(\xi), \sin(\xi)]$ to zero and we get the following nonlinear algebraic system :

$$\cos^{10}(\xi) : -60480A_4c^2 + 72A_4^2 - 72B_4^2,$$

$$\cos^9(\xi) : -20160A_3c^2 + 112A_3A_4 - 112B_4B_3,$$

$$\cos^9(\xi) \sin(\xi) : -60480B_4c^2 + 144A_4B_4,$$

$$\begin{aligned} \cos^8(\xi) : & -5040A_2c^2 + 195720A_4c^2 + 84A_2A_4 \\ & + 42A_3^2 - 128A_4^2 + \mu 840A_4 - 84B_2B_4 - 42B_3^2 + 170B_4^2, \end{aligned}$$

$$\cos^8(\xi) \sin(\xi) : -20160B_3c^2 + 112A_3B_4 + 112A_3B_4,$$

$$\begin{aligned} \cos^7(\xi) : & -720A_1c^2 + 60120A_3c^2 + 60A_1A_4 + 60A_2A_3 \\ & - 196A_3A_4 + \mu 360A_3 - 60B_1B_4 - 60B_2B_3 + 256B_3B_4, \end{aligned}$$

$$\begin{aligned} \cos^7(\xi) \sin(\xi) : & -5040B_2c^2 - 168000B_4c^2 \\ & + 84A_2B_4 + 84A_3B_3 + 84A_4B_2 - 226A_4B_4 + \mu 840B_4, \end{aligned}$$

$$\begin{aligned} \cos^6(\xi) : & 13560A_2c^2 - 232820A_4c^2 + 40A_0A_4 + 40A_1A_3 \\ & + 20A_2^2 - 144A_2A_4 + \mu 120A_2 - 72A_3^2 + 56A_4^2 - \mu 2080A_4 \\ & - 40B_1B_3 - 20B_2^2 + 184B_2B_4 + 92B_3^2 - 128B_4^2 + 20A_4, \end{aligned}$$

$$\begin{aligned} \cos^6(\xi) \sin(\xi) : & -720B_1c^2 + 50400B_3c^2 + 60A_1B_4 + 60A_2B_3 \\ & + 60A_3B_2 - 170A_3B_4 + 60A_4B_1 - 170A_4B_3 + \mu 360B_3, \end{aligned}$$

$$\begin{aligned} \cos^5(\xi) : & 1704A_1c^2 - 63660A_3c^2 + 24A_0A_3 + 24A_1A_2 \\ & - 100A_1A_4 + \mu 24A_1 - 100A_2A_3 + 84A_3A_4 - \mu 816A_3 \\ & - 24B_1B_2 + 124B_1B_4 + 124B_2B_3 - 184B_3B_4 + 12A_3, \end{aligned}$$

$$\begin{aligned} \cos^5(\xi) \sin(\xi) : & 11040B_2c^2 - 163160B_4c^2 + 40A_0B_4 \\ & + 40A_1B_3 + 40A_2B_2 - 122A_2B_4 + 40A_3B_1 - 122A_3B_3 - 122A_4B_2 \\ & + 84A_4B_4 + \mu 120B_2 - \mu 1720B_4 + 20B_4, \end{aligned}$$

$$\begin{aligned}
\cos^4(\xi) &: -12342A_2c^2 + 121280A_4c^2 + 12A_0A_2 - 64A_0A_4 + 6A_1^2 - 64A_1A_3 \\
&- 32A_2^2 + 60A_2A_4 - \mu 240A_2 + 30A_3^2 + \mu 1696A_4 - 6B_1^2 + 76B_1B_3 + 38B_2^2 \\
&- 124B_2B_4 - 62B_3^2 + 30B_4^2 + 6A_2 - 32A_4, \\
\cos^4(\xi) \sin(\xi) &: 1344B_1c^2 - 41832B_3c^2 + 24A_0B_3 + 24A_1B_2 - 82A_1B_4 + 12B_3 \\
&+ 24A_2B_1 - 82A_2B_3 - 82A_3B_2 + 60A_3B_4 - 82A_4B_1 + 60A_4B_3 + \mu 24B_1 - \mu 648B_3, \\
\cos^3(\xi) &: -1274A_1c^2 + 27522A_3c^2 + 4A_0A_1 - 36A_0A_3 - 36A_1A_2 + 2A_1 - 18A_3 \\
&+ 40A_1A_4 - \mu 40A_1 + 40A_2A_3 + \mu 576A_3 + 40B_1B_2 - 76B_1B_4 - 76B_2B_3 + 40B_3B_4, \\
\cos^3(\xi) \sin(\xi) &: -7452B_2c^2 + 63091B_4c^2 + 12A_0B_2 - 50A_0B_4 + \mu 1061B_4 + 6B_2 - 25B_4 \\
&+ 12A_1B_1 - 50A_1B_3 - 50A_2B_2 + 40A_2B_4 - 50A_3B_1 + 40A_3B_3 + 40A_4B_2 - \mu 180B_2, \\
\cos^2(\xi) &: 4112A_2c^2 - 24684A_4c^2 - 16A_0A_2 + 24A_0A_4 - 8A_1^2 + 24A_1A_3 + 12A_2^2 \\
&+ \mu 136A_2 - \mu 480A_4 + 8B_1^2 - 40B_1B_3 - 20B_2^2 - 24B_2B_4 + 12B_3^2 - 8A_2 + 12A_4, \\
\cos^2(\xi) \sin(\xi) &: -692B_1c^2 + 12283B_3c^2 + 4A_0B_1 - 26A_0B_3 - 26A_1B_2 - 13B_3 \\
&+ 24A_1B_4 - 26A_2B_1 + 24A_2B_3 + 24A_3B_2 + 24A_4B_1 - \mu 28B_1 + \mu 317B_3 + 2B_1, \\
\cos(\xi) &: 290A_1c^2 - 3822A_3c^2 - 4A_0A_1 + 12A_0A_3 + 12A_1A_2 \\
&+ \mu 16A_1 - \mu 120A_3 - 16B_1B_2 + 12B_1B_4 + 12B_2B_3 - 2A_1 + 6A_3, \\
\cos(\xi) \sin(\xi) &: 1451B_2c^2 - 7452B_4c^2 - 10A_0B_2 + 12A_0B_4 - 10A_1B_1 + 12A_1B_3 \\
&+ 12A_2B_2 + 12A_3B_1 + \mu 61B_2 - \mu 180B_4 - 5B_2 + 6B_4, \\
\sin(\xi) &: 67B_1c^2 - 692B_3c^2 - 2A_0B_1 + 4A_0B_3 + 4A_1B_2 \\
&+ 4A_2B_1 + \mu 5B_1 - \mu 28B_3 - B_1 + 2B_3, \\
\text{constant} &: -290A_2c^2 + 984A_4c^2 + 4A_0A_2 + 2A_1^2 - \mu 16A_2 \\
&+ \mu 24A_4 - 2B_1^2 + 4B_1B_3 + 2B_2^2 + 2A_2.
\end{aligned}$$

(3.4)

We solve the set of nonlinear algebraic equations with the help of Maple, the solutions of these algebraic equations are found to be in the following.

Case 1:

$$\begin{aligned} A_0 &= \frac{1105}{2}c^2 - \frac{1}{2}, & A_1 &= 0, & A_2 &= -1680c^2, & A_3 &= 0, & A_4 &= 840c^2, \\ B_1 &= 0, & B_2 &= 0, & B_3 &= 0, & B_4 &= 0, & \mu &= 51c^2, & c &= \frac{1}{4}. \end{aligned} \quad (3.5)$$

If we replace these results into (3.3) and insert the result into the transformation (2.2), we acquire the exact solitary wave solution of equation:

$$u_1(x, t) = -\frac{1}{2} \frac{575c^2 \cosh^4(x - ct) + \cosh^4(x - ct) - 1680c^2}{\cosh^4(x - ct)}. \quad (3.6)$$




FIGURE 1. Figure 1 shows that the 3D and 2D soliton solution (3.6) for $c = \frac{1}{4}$, $-5 < x < 5$, $-5 < t < 5$.

Case 2:

$$\begin{aligned} A_0 &= -\frac{1247}{2}c^2 - 1368c^2\left(-\frac{1}{2} + \frac{1}{62}I\sqrt{31}\right) - \frac{1}{2}, \\ A_1 &= 0, & A_2 &= 1680c^2\left(-\frac{1}{2} + \frac{1}{62}I\sqrt{31}\right), & A_3 &= 0, \\ A_4 &= 840c^2, & B_1 &= 0, & B_2 &= 0, & B_3 &= 0, \\ B_4 &= 0, & \mu &= -105c^2 - 156c^2\left(-\frac{1}{2} + \frac{1}{62}I\sqrt{31}\right). \end{aligned} \quad (3.7)$$

If we replace these results into (3.3), we get the following solution:

$$\begin{aligned} u_2(x, t) &= \frac{1}{62}I \left(\frac{312\sqrt{31}c^2 \cosh^4(x - ct) - 1680\sqrt{31}c^2 \cosh^2(x - ct) + 375c^2 \cosh^4(x - ct)}{\cosh^4(x - ct)} \right) \\ &+ \left(\frac{52080c^2 - c^2 52080 \cosh^2(x - ct) - 31 \cosh^4(x - ct)}{\cosh^4(x - ct)} \right). \end{aligned} \quad (3.8)$$




FIGURE 2. Figure 2 The 3D and 2D surfaces of solution (3.8) with respectively real part and imaginary part .

Case 3:

$$\begin{aligned}
 A_0 &= \frac{385}{2}c^2 - \frac{1}{2}, & A_1 &= 0, & A_2 &= -630c^2, & A_3 &= 0, & A_4 &= 420c^2, \\
 B_2 &= -420Ic^2, & B_3 &= 0, & B_4 &= 420Ic^2, & \mu &= 12c^2, & B_1 &= 0.
 \end{aligned}
 \tag{3.9}$$

Similar to the previous case, we get the following solitary wave solution:

$$u_3(x, t) = -\frac{1}{2} \left(\frac{35c^2 \cosh^4(x - ct) + \cosh^4(x - ct)}{\cosh^4(x - ct)} \right) \tag{3.10}$$

$$+ \frac{420 \cosh^2(x - ct)c^2 - 840c^2}{\cosh^4(x - ct)} + I \left(\frac{840c^2 \sinh(x - ct)}{\cosh^4(x - ct)} \right) \tag{3.11}$$




FIGURE 3. The 3D and 2D surfaces of solution (3.1) with respectively real part and imaginary part.

Case 4:

$$\begin{aligned}
 A_0 &= -\frac{381}{4}\left(-\frac{3}{4} + \frac{1}{124}I\sqrt{31}\right)\sqrt{-\frac{1}{11025}B_4^2} - \frac{353}{8}\sqrt{-\frac{1}{11025}B_4^2} - \frac{1}{2}, \\
 A_1 &= 0, \quad A_3 = 0, \quad B_1 = 0, \\
 A_2 &= \frac{105}{2}\left(-\frac{5}{2} + \frac{1}{62}I\sqrt{31}\right)\sqrt{-\frac{1}{11025}B_4^2}, \\
 A_4 &= 105\sqrt{-\frac{1}{11025}B_4^2}, \quad B_2 = B_4\left(-\frac{3}{4} + \frac{1}{124}I\sqrt{31}\right), \\
 B_3 &= 0, \quad c = \frac{1}{2}\left(-\frac{1}{11025}B_4^2\right)^{\frac{1}{4}}, \\
 \mu &= -\frac{3}{2}\sqrt{-\frac{1}{11025}B_4^2}\left(\frac{5}{4} + \frac{13}{124}I\sqrt{31}\right), \quad B_4 = 1.
 \end{aligned} \tag{3.12}$$




FIGURE 4. The 3D and 2D surfaces of solution (3.13) with respectively real part and imaginary part.

Similar to the previous case, we get the following solitary wave solution:

$$\begin{aligned}
 u_4(\xi) = & \left(\frac{1}{52080} \frac{39I\sqrt{-B_4^2}\sqrt{31}\cosh^4(\xi) + 420IB_4\sinh(\xi)\sqrt{31}\cosh^2(\xi) + 527\sqrt{-B_4^2}\cosh^4(\xi)}{\cosh^4(\xi)} \right) \\
 & - \left(\frac{420I\sqrt{-31B_4^2}\cosh^2(\xi)}{\cosh^4(\xi)} \right) \\
 & + \left(\frac{1}{52080} \frac{13020B_4\sinh(\xi)\cosh^2(\xi) - 26040\cosh^4(\xi) - 39060\cosh^2(\xi)\sqrt{B_4^2}}{\cosh^4(\xi)} \right) \\
 & - \left(\frac{52080B_4\sinh(\xi) + 52080\sqrt{B_4^2}}{\cosh^4(\xi)} \right)
 \end{aligned} \tag{3.13}$$

3.2. Kuramoto-Sivashinsky Equation. Secondly, we consider Kuramoto-Sivashinsky equation:

$$u_t + \alpha uu_x + bu_{xx} + ku_{xxx} = 0, \quad (3.14)$$

In this subsection we are going to sustain the exact traveling wave solutions of the Kuramoto-Sivashinsky equation by using of the sine-Gordon method.

We look for the solutions of the Kuramoto-Sivashinsky equation by the method described in past section. We operate the travelling wave transformation of the following form,

$$u(x, t) = u(\xi), \quad \xi = x - ct, \quad (3.15)$$

Eq.(3.14) is reduced into the following ordinary differential equations (ODE)

$$-cu' + \alpha uu' + bu'' + ku'''' = 0, \quad (3.16)$$

where prime denotes differentiation with respect to ξ . If we balance between the highest order nonlinear derivative u'''' and nonlinear term of the highest degree uu' in Eq.(3.16), we obtain $n = 3$.

Hence, the expected solution takes the form

$$u(\xi) = A_0 + B_1 \operatorname{sech}(\xi) + B_2 \operatorname{sech}(\xi) \tanh(\xi) + B_3 \operatorname{sech}(\xi) \tanh^2(\xi) + A_1 \tanh(\xi) + A_2 \tanh^2(\xi) + A_3 \tanh^3(\xi), \quad (3.17)$$

where $u = u(\xi)$ satisfies Eq.(2.5) in which $A_0, A_1, A_2, A_3, B_1, B_2, B_3$ are undetermined constants.

We substitute Eq.(3.17) into Eq.(3.16) secondly utilize Eq.(2.5) finally we equate all coefficients of the functions $[\cos(\xi), \sin(\xi)]$ to zero and we get:

$$\cos^7(\xi) : -3A_3^2 a + 3B_2^3 a + 360A_3 k,$$

$$\cos^6(\xi) : -5A_2 A_3 a + 5B_2 B_3 a + 120A_2 k,$$

$$\cos^6(\xi) \sin(\xi) : -6A_3 B_3 a + 360B_3 k,$$

$$\begin{aligned} \cos^5(\xi) : & -4A_1 A_3 a - 2A_2^2 a + 3A_3^2 a \\ & + 4B_1 B_3 a + 2B_2^2 a - 5B_3^2 a + 24A_1 k + 12A_3 b - 816A_3 k, \end{aligned}$$

$$\begin{aligned}
\cos^5(\xi) \sin(\xi) &: -5A_2B_3a - 5A_3B_2a + 120B_2k, \\
\cos^4(\xi) &: -3A_0A_3a - 3A_1A_2a + 5A_2A_3a + 3B_1B_2a - 8B_2B_3a + 6A_2b - 240A_2k + 3A_3c, \\
\cos^4(\xi) \sin(\xi) &: -4A_1B_3a - 4A_2B_2a - 4A_3B_1a + 5A_3B_3a \\
&+ 24B_1k + 12B_3b - 648B_3k, \\
\cos^3(\xi) &: -2A_0A_2a - A_1^2a + 4A_1A_3a + 2A_2^2a + B_1^2a - 6B_1B_3a \\
&- 3B_2^2a + 2B_3^2a + 2A_1b - 40A_1k + 2A_2c - 18A_3b + 576A_3k, \\
\cos^3(\xi) \sin(\xi) &: -3A_0B_3a - 3A_1B_2a - 3A_2B_1a + 4A_2B_3a \\
&+ 4A_3B_2a + 6B_2b - 180B_2k + 3B_3c, \\
\cos^2(\xi) &: -A_0A_1a + 3A_0A_3a + 3A_1A_2a - 4B_1B_2a + 3B_2B_3a + A_1c \\
&- 8A_2b + 136A_2k - 3A_3c, \\
\cos^2(\xi) \sin(\xi) &: -2A_0B_2a - 2A_1B_1a + 3A_1B_3a \\
&+ 3A_2B_2a + 3A_3B_1a + 2B_1b - 28B_1k + 2B_2c \\
&- 13B_3b + 317B_3k, \\
\cos(\xi) &: 2A_0A_2a + A_1^2a - B_1^2a + 2B_1B_3a + B_2^2a - 2A_1b \\
&+ 16A_1k - 2A_2c + 6A_3b - 120A_3k, \\
\cos(\xi) \sin(\xi) &: -A_0B_1a + 2A_0B_3a + 2A_1B_2a + 2A_2B_1a + B_1c \\
&- 5B_2b + 61B_2k - 2B_3c, \\
\sin(\xi) &: A_0B_2a + A_1B_1a - B_1b + 5B_1k - B_2c + 2B_3b - 28B_3k, \\
\text{constant} &: A_0A_1a + B_1B_2a - A_1c + 2A_2b - 16A_2k.
\end{aligned} \tag{3.19}$$

We solved the set of nonlinear algebraic equations with the help of Maple, the solutions of these algebraic equations are found to be in the following.

Set 1:

$$A_0 = A_0, A_1 = -\frac{9}{11}A_3, A_2 = 0, , \quad (3.20)$$

$$B_1 = 0, B_2 = 0, B_3 = 0, \quad (3.21)$$

$$b = \frac{19}{330}A_3a, \quad c = A_0a, \quad (3.22)$$

$$k = \frac{1}{120}A_3a. \quad (3.23)$$

Set 2:

$$A_1 = IB_1, \quad A_2 = 0, \quad A_3 = 0, \quad (3.24)$$

$$B_2 = 0, \quad B_3 = 0, \quad b = IaB_1, \quad (3.25)$$

$$c = A_0a, \quad k = 0. \quad (3.26)$$

Set 3:

$$\begin{aligned} A_1 &= -\frac{3}{2}IB_3, \quad A_2 = 0, \quad A_3 = IB_3, \\ B_1 &= -B_3, \quad B_2 = 0, \\ b &= -\frac{19}{60}IaB_3, \quad c = A_0a, \\ k &= \frac{1}{60}IaB_3. \end{aligned} \quad (3.27)$$

Set 4:

$$\begin{aligned} A_1 &= -\frac{21}{22}IB_3, \quad A_2 = 0, \quad A_3 = IB_3, \\ B_1 &= -\frac{5}{11}B_3, \quad B_2 = 0, \\ b &= \frac{19}{660}IaB_3, \quad c = A_0a, \\ k &= \frac{1}{60}IaB. \end{aligned} \quad (3.28)$$

If we substitute these results into (3.3) we attain the exact travelling wave solutions of equation:

$$u_1(x, t) = A_0 - \frac{9}{11}A_3 \tanh(x - A_0at) + A_3 \tanh^3(x - A_0at). \quad (3.29)$$




FIGURE 5. 3D and 2D soliton solution (3.29) for $A_0 = 1/2$, $a = 1/2$, $A_3 = 1$.

$$u_2(x, t) = \frac{IB_1 \sinh(x - A_0at) + A_0 \cosh(x - A_0at) + B_1}{\cosh(x - A_0at)}. \quad (3.30)$$




FIGURE 6. For $A_0 = 1/2$, $a = 1/2$, $B_1 = 1$ The 3D and 2D surfaces of solution (3.30) with respectively real part and imaginary part.

$$u_3(x, t) = \frac{1}{2} \left(\frac{2A_0 \cosh^3(x - A_0at) - -2B_3}{\cosh^3(x - A_0at)} \right) \quad (3.31)$$

$$-I \frac{B_3 \sinh(x - A_0at) \cosh^2(x - A_0at)}{\cosh^3(x - A_0at)} \quad (3.32)$$

$$+I \frac{+2B_3 \sinh(x - A_0at)}{\cosh^3(x - A_0at)} \quad (3.33)$$




FIGURE 7. For $A_0 = 1/2$, $a = 1/2$, $B_3 = 1$. The 3D and 2D surfaces of solution (3.33) with respectively real part and imanigar part.

$$u_4(x, t) = \frac{1}{22} \left(\frac{IB_3 \sinh(x - A_0at) \cosh^2(x - A_0at) + 22A_0 \cosh^3(x - A_0at)}{\cosh^3(x - A_0at)} \right) + \frac{1}{22} \left(\frac{-22IB_3 \sinh(x - A_0at) + 12B_3 \cosh^2(x - A_0at) - 22B_3}{\cosh^3(x - A_0at)} \right). \quad (3.34)$$




FIGURE 8. For $A_0 = 1/2$, $a = 1/2$, $B_3 = 1$.
 The 3D and 2D surfaces of solution(3.34)
 with respectively real part and imaginary part.

3.3. Sawada-Kotera Equation. As a last application of the related method, we consider Sawada-Kotera equation and is given as follows:

$$u_t + [63u^4 + 63(2u^2u_{xx} + uu_x^2) + 21(uu_{xxxx} + u_{xx}^2 + u_xu_{xxx}) + u_{xxxxx}]_x = 0, \quad (3.35)$$

By substituting transformation $u(x, t) = u(\xi)$, $\xi = x - ct$, Eq.(3.35) is transformed into an one dimension form

$$-cu' + 252u^3u' + 378uu''u' + 126u^2u''' + 63(u')^3 + 42u'u'''' + 21uu'''' + 63u''u''' + u^{vii} = 0, \quad (3.36)$$

where prime denotes differentiation with respect to ξ . The balance between u^{vii} and u^3u' gives $n = 2$.

Thereby, the exact solution of the proposed equation has the form,

$$u(\xi) = A_0 + B_1 \operatorname{sech}(\xi) + B_2 \operatorname{sech}(\xi) \tanh(\xi) + A_1 \tanh(\xi) + A_2 \tanh^2(\xi), \quad (3.37)$$

where $u = u(\xi)$ satisfies Eq.(2.5) in which A_0, A_1, A_2, B_1, B_2 are undetermined constants.

As in other applications, the algebraic equation system construct and when it is solved, the following solution sets are obtained.

Set 1:

$$A_1 = 0, \quad A_2 = -2, \quad B_1 = 0, \quad (3.38)$$

$$B_2 = 0, \quad c = 252A_0^3 - 1008A_0^2 + 1344A_0 - 608. \quad (3.39)$$

Set 2:

$$A_1 = 0, \quad A_2 = -1, \quad B_1 = 0, \quad (3.40)$$

$$B_2 = I, \quad c = 252A_0^3 - 630A_0^2 + 525A_0 - 146. \quad (3.41)$$

Set 3:

$$A_0 = \frac{8}{3}, \quad A_1 = 0, \quad A_2 = -4, \quad (3.42)$$

$$B_1 = 0, \quad B_2 = 0, \quad c = -\frac{256}{3}. \quad (3.43)$$

Set 4:

$$A_0 = \frac{5}{3}, \quad A_1 = 0, \quad A_2 = -2, \quad (3.44)$$

$$B_1 = 0, \quad B_2 = 2I, \quad c = -\frac{4}{3}. \quad (3.45)$$

If we substitute these results into (3.3) we attain the exact travelling wave solutions of equation:

$$u_1(\xi) = \frac{A_0 \cosh^2(\xi) - 2 \cosh^2(\xi) + 2}{\cosh^2(\xi)}. \quad (3.46)$$

$$u_2(\xi) = \frac{I \sinh(\xi) + A_0 \cosh^2(\xi) + 1 - \cosh^2(\xi)}{\cosh^2(\xi)} \quad (3.47)$$

$$u_3(\xi) = -\frac{4}{3} \left(\frac{\cosh^2(\xi) - 3}{\cosh^2(\xi)} \right). \quad (3.48)$$

$$u_4(\xi) = \frac{1}{3} \left(\frac{6I \sinh(\xi) - \cosh^2(\xi) + 6}{\cosh^2(\xi)} \right). \quad (3.49)$$




FIGURE 9. For $A_0 = 1$.
 The 3D and 2D surfaces of solution(3.47)
 with respectively real part and imaginary part.

4. CONCLUSION

In this article, we studied on acquiring the exact travelling wave solutions of (1+1)-dimensional nonlinear equations by introducing appropriate transformations and applying the sine-gordon expansion method. We employed this method to find the solutions of Higher-Order Boussinesq, Kuramoto-Sivashinsky and Sawada-Kotera Equations. We obtained the sets of nonlinear equations that can be solved by using Maple software.

The purpose of this study was constructed on reliable, beneficial and consistent treatment for the analysis of the above-mentioned equations. We can say that the sine-gordon expansion method is a very powerful and efficient mathematical tool to solve nonlinear equations. Furthermore, we believe that this method is also useful for a variety of other nonlinear equations that represent in mathematical, physics and other nonlinear sciences. The solutions acquired may be very beneficial in order to understand the mechanism of the intricate nonlinear physical phenomena of referred equations. The conclusions obtained will be useful for guiding future research works in the appropriate fields.

Declaration of Competing Interest

Authors have no interests to declare.

REFERENCES

- [1] Zhang, J., Jiang, F., Zhao, X.: An improved (G0/G)-expansion method for solving nonlinear evolution equations. *Int. J. Math.* 87, 1716–1725 (2010)
- [2] Seadawy, A.R., Lu, D., Khater, M.M.A.: Solitary wave solutions for the generalized Zakharov–Kuznetsov– Benjamin–Bona–Mahony nonlinear evolution equation. *J. Ocean Eng. Sci.* 2(2), 137–142 (2017b)
- [3] Naher, Hasibun, and Farah Aini Abdullah. "New Traveling Wave Solutions by the Extended Generalized Riccati Equation Mapping Method of the-Dimensional Evolution Equation." *Journal of Applied Mathematics* 2012 (2012).
- [4] Bulut, H., Yel, G., Baskonus, H.M.: Novel structure to the coupled nonlinear Maccari's system by using modified trial equation. *Adv. Math. Models Appl.* 2(1), 14–19 (2017)
- [5] Bulut, H., Baskonus, H.M.: A study on the numerical solution of the third-order dispersive equations with homotopy perturbation method. *e-J. New World Sci. Acad.* 5(1), 18 (2010)
- [6] Hosseini, K., M. Mirzazadeh, and J. F. Gómez-Aguilar. "Soliton solutions of the Sasa–Satsuma equation in the monomode optical fibers including the beta-derivatives." *Optik* 224 (2020): 165425.
- [7] Hosseini, K., et al. "A high-order nonlinear Schrödinger equation with the weak non-local nonlinearity and its optical solitons." *Results in Physics* 23 (2021): 104035.
- [8] Hosseini, Kamyar, Mashaallah Matinfar, and Mohammad Mirzazadeh. "Soliton solutions of high-order nonlinear Schrödinger equations with different laws of nonlinearities." *Regular and Chaotic Dynamics* 26.1 (2021): 105-112.
- [9] Hosseini, K., et al. "An integrable (2+ 1)-dimensional nonlinear Schrödinger system and its optical soliton solutions." *Optik* 229 (2021): 166247.
- [10] Hosseini, Kamyar, et al. "The (2+ 1)-dimensional Heisenberg ferromagnetic spin chain equation: its solitons and Jacobi elliptic function solutions." *The European Physical Journal Plus* 136.2 (2021): 1-9.
- [11] Yaşar, E., and I. B. Giresunlu. "Lie symmetry reductions, exact solutions and conservation laws of the third order variant Boussinesq system." *Acta Physica Polonica A* 128.3 (2015): 252-255.
- [12] San, Sait, and Emrullah Yaşar. "On the conservation laws and exact solutions of a modified Hunter-Saxton equation." *Advances in Mathematical Physics* 2014 (2014).
- [13] Kaplan, Melike, Sait San, and Ahmet Bekir. "On the exact solutions and conservation laws to the Benjamin-Ono equation." *Journal of Applied Analysis & Computation* 8.1 (2018): 1-9.
- [14] Schneider, Guido, and C. Eugene Wayne. "Kawahara dynamics in dispersive media." *Physica D: Nonlinear Phenomena* 152 (2001): 384-394.
- [15] Gözükızıl, Ömer Faruk, and Şamil Akça ğıl. "Travelling wave solutions to the Benney-Luke and the higher-order improved Boussinesq equations of Sobolev type." *Abstract and applied analysis*. Vol. 2012. Hindawi, 2012.
- [16] Hao, Jianghao, and Aiyuan Gao. "Blow-up for the sixth-order multidimensional generalized Boussinesq equation with arbitrarily high initial energy." *Boundary Value Problems* 2019.1 (2019): 1-13.
- [17] Wang, Shubin, and Hongxia Xue. "Global solution for a generalized Boussinesq equation." *Applied mathematics and computation* 204.1 (2008): 130-136.

- [18] C. Li, G. Chen and S. Zhao, Exact Travelling wave solutions to the generalized Kuramoto-Sivashinsky equation, *Latin American Applied Research*, 34, 64-68 (2004).
- [19] Kuramoto, Y., and Tsuzuki, T., 1976, "Persistent Propagation of Concentration Waves in Dissipative Media far from Thermal Equilibrium," *Prog. Theor. Phys.*, 55, pp. 356–369.
- [20] LaQuey, R., Mahajan, S., Rutherford, P., and Tang, W., 1975, "Nonlinear Saturation of the Trapped-Ion Mode," *Phys. Rev. Lett.*, 34, pp. 391–394.
- [21] Sivashinsky, G., 1977, "Nonlinear Analysis of Hydrodynamic Instability in Laminar Flames—I. Derivation of Basic Equations," *Acta Astron.*, 4, pp. 1177–1206.
- [22] Kurulay, Muhammet, Aydin Secer, and Mehmet Ali Akinlar. "A new approximate analytical solution of Kuramoto-Sivashinsky equation using homotopy analysis method." *Applied Mathematics & Information Sciences* 7.1 (2013): 267-271.
- [23] Khater, A. H., and R. S. Temsah. "Numerical solutions of the generalized Kuramoto-Sivashinsky equation by Chebyshev spectral collocation methods." *Computers & Mathematics with Applications* 56.6 (2008): 1465-1472.
- [24] Aljahdaly, Noufe H., Aly R. Seadawy, and Wafaa A. Albarakati. "Applications of dispersive analytical wave solutions of nonlinear seventh order Lax and Kaup-Kupershmidt dynamical wave equations." *Results in Physics* 14 (2019): 102372.
- [25] Seadawy, A., New exact solutions for the KdV equation with higher order nonlinearity by using the variational method, *Computers and Mathematics with Applications*, 62 (2011), 10, pp. 3741–3755.
- [26] Kabir, M., Modified Kudryashov method for finding exact solitary wave solutions of higher-order nonlinear equations, *Mathematical methods in the Applied Sciences*, 34 (2011), 2, pp. 213–219.
- [27] Ganji, D., et. al., New exact solutions for seventh-order Sawada-Kotera-Ito, Lax and Kaup-Kupershmidt equations using exp-function method, *Mathematical Methods in the Applied Sciences*, 33 (2010), 2, pp.167–176.
- [28] Feng, J., New traveling wave solutions to the seventh-order Sawada-Kotera equation, *Journal of applied mathematics & informatics*, 28 (2010), 5-6, pp.1431–1437.
- [29] Salas, A. H., et. al., Computing exact solutions to a generalized Lax-Sawada-Kotera-Ito seventh-order KdV equation, *Mathematical Problems in Engineering*, 2010 (2010), pp. 1–7.
- [30] Jafari, Hossein, et al. "Application of He's variational iteration method for solving seventh order Sawada-Kotera equations." *Applied Mathematical Sciences* 2.9-12 (2008): 471-477.
- [31] Wazwaz, Abdul-Majid. "The Hirota's direct method and the tanh-coth method for multiple-soliton solutions of the Sawada-Kotera-Ito seventh-order equation." *Applied Mathematics and Computation* 199.1 (2008): 133-138.
- [32] Shen, Yu-Jia, et al. "Bell-polynomial approach applied to the seventh-order Sawada-Kotera-Ito equation." *Applied Mathematics and Computation* 227 (2014): 502-508.
- [33] Korkmaz, A., Hepson, O. E., Hosseini, K., Rezazadeh, H., & Eslami, M. (2020). Sine-Gordon expansion method for exact solutions to conformable time fractional equations in RLW-class. *Journal of King Saud University-Science*, 32(1), 567-574.
- [34] Yamgoué, Serge Bruno, Guy Roger Deffo, and François Beceau Pelap. "A new rational sine-Gordon expansion method and its application to nonlinear wave equations arising in mathematical physics." *The European Physical Journal Plus* 134.8 (2019): 1-15.

- [35] Kumar, D., Kamyar Hosseini, and F. Samadani. "The sine-Gordon expansion method to look for the traveling wave solutions of the Tzitzé ica type equations in nonlinear optics." *Optik* 149 (2017): 439-446.
- [36] Korkmaz, Alper, et al. "Sine-Gordon expansion method for exact solutions to conformable time fractional equations in RLW-class." *Journal of King Saud University-Science* 32.1 (2020): 567-574.
- [37] Ali, Khaled K., Abdul-Majid Wazwaz, and M. S. Osman. "Optical soliton solutions to the generalized nonautonomous nonlinear Schrödinger equations in optical fibers via the sine-Gordon expansion method." *Optik* 208 (2020): 164132.